

**From:** [Kelley Jorgensen](#)  
**To:** [Jenna Kay](#); [Brent Davis](#)  
**Cc:** [David Morgan](#); [Chris Watson](#); [Kelley Jorgensen](#)  
**Subject:** [Contains External Hyperlinks] RE: Zip shapes for delivery  
**Date:** Thursday, December 5, 2019 4:26:11 PM  
**Attachments:** [image001.png](#)  
[DRAFT Plas Newydd Wapato Valley OHWM Determination 12.5.2019 redux.pdf](#)

---

**CAUTION:** This email originated from outside of Clark County. Do not click links or open attachments unless you recognize the sender and know the content is safe.

Jenna and Brent,

Please find attached our draft report documenting the OHWM determination for Wapato Valley Mitigation and Conservation Bank and Plas Newydd Farm. We do not propose any change to Allen Creek (or Allen Canyon Creek) or Lake Rosannah at this time, so those waterbodies are not detailed in the report.

Please confirm receipt of this PDF, and don't hesitate to contact me about the report, or Chris Watson about the GIS shapefiles.

Thank you for the opportunity to provide input to the process.

Kelley



KELLEY JORGENSEN  
President of Conservation  
she | her | hers

T 360.857.4087 C 971.285.6874  
E [kjorgensen@pnfarm.com](mailto:kjorgensen@pnfarm.com)

PO Box 428  
Ridgefield, WA 98642 | [www.wapato-valley.com](http://www.wapato-valley.com)

---

**From:** Jenna Kay <[Jenna.Kay@clark.wa.gov](mailto:Jenna.Kay@clark.wa.gov)>  
**Sent:** Thursday, December 5, 2019 10:36 AM  
**To:** Kelley Jorgensen <[kjorgensen@pnfarm.com](mailto:kjorgensen@pnfarm.com)>; Brent Davis <[Brent.Davis@clark.wa.gov](mailto:Brent.Davis@clark.wa.gov)>  
**Subject:** RE: Zip shapes for delivery

Thanks Kelley. We will follow-up once we have a chance to review and look forward to receiving the additional document.

Jenna

---

**From:** Kelley Jorgensen [[kjorgensen@pnfarm.com](mailto:kjorgensen@pnfarm.com)]  
**Sent:** Wednesday, December 04, 2019 4:58 PM  
**To:** Brent Davis; Jenna Kay  
**Subject:** [Contains External Hyperlinks] FW: Zip shapes for delivery

**CAUTION:** This email originated from outside of Clark County. Do not click links or open attachments unless you recognize the sender and know the content is safe.

Hi Brent and Jenna,

Please find attached the wetland rating unit, OHWM and 100-year flood GIS shapefiles for the Plas

Newydd property.

These are the locations we are proposing updates for the Shoreline Master Plan process.

OHWL delineation technical memo to follow under separate cover.

Please don't hesitate to contact me or Chris Watson if you have any questions.

Thank you for your time and consideration,

Kelley

---

**KELLEY JORGENSEN** » *President of Conservation*


*she | her | hers*

T 360.857.4087 C 971.285.6874 E [kjorgensen@pnfarm.com](mailto:kjorgensen@pnfarm.com)

PO Box 428 | Ridgefield, WA 98642 | [www.wapato-valley.com](http://www.wapato-valley.com)

Please find attached GIS shapefiles:

- 100yr flood zone on the PN Farm property based on the USACE 1% exceedance value at Columbia RM 87 of 26.54' NAVD 88.



- Wetland rating units on the Wapato Mitigation and Conservation Bank provided by CEG.
- OHWM from determination.

**CHRIS WATSON** » *GIS Manager/Project*

*Manager/Geologist*

*he | him | his*

---

T 360.857.4040 C 503.332.7231 E [cwatson@pnfarm.com](mailto:cwatson@pnfarm.com)

PO Box 428 | Ridgefield, WA 98642 | [www.wapato-valley.com](http://www.wapato-valley.com)

This e-mail and related attachments and any response may be subject to public disclosure under state law.

# DETERMINING THE ORDINARY HIGH WATER MARK FOR THE WAPATO VALLEY MITIGATION AND CONSERVATION BANK AND PLAS NEWYDD FARM

DECEMBER 2019



PLAS NEWYDD FARM  
EST. 1941  
CONSERVATION PROGRAM

Prepared by  
Plas Newydd, LLC  
Conservation Program  
Ridgefield, Washington

Prepared for  
Clark County, WA & Washington IRT

**Cover Photos showing diversity of shoreline conditions, clockwise from upper left:**

1. Native basalt outcrop with moss scour line, Gee Creek backwater south of the Narrows Levee, Gee Creek approx. RM 2.33
2. Columbia River shoreline with flattened emergent vegetation, approx. RM 87.1
3. Lewis River shoreline with sandy bank wrack line, approx. RM 0.1
4. Gee Creek shoreline vegetation transition, approx. RM 1.95

**Suggested citation:**

Plas Newydd, Inc. 2019. Determining the Ordinary High Water Mark for the Wapato Valley Mitigation and Conservation Bank. Ridgefield, Washington.

# TABLE OF CONTENTS

**1 INTRODUCTION ..... 1**

**2 METHODS..... 2**

**3 FIELD ASSESSMENT..... 4**

    3.1 COLUMBIA RIVER OHWM..... 5

    3.2 LEWIS RIVER OHWM..... 6

    3.3 GEE CREEK OHWM ..... 7

    3.4 LANCASTER LAKE OHWM ..... 7

**4 HYDROLOGIC ASSESSMENT METHODS..... 9**

**5 STREAM HYDROLOGIC ASSESSMENT ..... 10**

    5.1 STEP 1 AND 2: USE GAGE DATA TO APPROXIMATE UPPER AND LOWER EXTREMES FOR  
OHWM FLOWS AND CORRELATE TO STAGE ..... 10

        5.1.1 Generate the upper bookends by estimating the two-year peak  
and minimum peak flow..... 10

        5.1.2 Refine the Range..... 12

        5.1.3 Step 3: Compare recent events to OHWM bookends ..... 13

        5.1.4 Stream assessment conclusions..... 13

**6 TIDAL HYDROLOGIC ASSESSMENT..... 14**

    6.1 STEPS 1–3: LOCATE AN APPROPRIATE STATION AND IDENTIFY TIDAL DATUMS ..... 14

    6.2 TIDAL ASSESSMENT CONCLUSIONS ..... 15

**7 CONCLUSIONS..... 15**

**8 REFERENCES..... 16**

**List of Tables**

Table 1. Waterbodies and Shoreline Areas included in Delineation of OHWM4

Table 2. Plant Distribution across Columbia River OHWM Gradient ..... 6

Table 3. Plant Distribution across Lewis River OHWM Gradient..... 7

Table 4. Plant Distribution across Gee Creek OHWM Gradient ..... 7

Table 5. Plant Distribution across Lancaster Lake OHWM Gradient ..... 8

Table 6. Maximum peak annual discharge data 1958–2017 Lewis River (aka  
“spreadsheet method”). ..... 10

Table 7. Number of times 16,400 cfs was exceeded in each year 2002–2017.  
..... 13

Table 8. Local Datum Comparisons to MHHW at St. Helens Tidal Station..... 14

Table 9. Hydrologic assessment “bookend” OHWM elevation ranges..... 15

Table 10. OHWM Results for Plas Newydd Farm/Wapato Valley in NAVD88 .. 15

**List of Figures**

Figure 1. Property Overview and Vicinity

Figure 2. OHWM Elevations

Figure 3. Location of the nearest tidal station and stream gage to Wapato Valley.

Figure 2. Hydrograph of the maximum peak annual discharge data for the Lewis River 2-year and 1.01-year peak flows depicted.

Figure 3. Stage for the determined flow range values plotted on aligned discharge and stage graphs.

Figure 4. Daily discharge plotted with refined OHWM bookend limits from refined analysis.

**Attachments**

Attachment A. Field Data Sheets and Maps

Attachment B. Species and Common Names of Plants

## 1 INTRODUCTION

Plas Newydd LLC proposes to construct and operate a wetland mitigation and habitat conservation bank, the Wapato Valley Mitigation and Conservation Bank (Wapato Valley or Bank), on privately owned land known as Plas Newydd Farm (PN Farm). The purpose of the Bank is to generate mitigation credits for projects that will have an adverse impact on the aquatic and adjacent terrestrial environment, and that need to compensate for those impacts as a condition of their permits or other regulatory requirements resulting from project impacts. The Bank also serves a critical purpose to conserve an important and rare landscape and the ecological processes that shape and define it, as well as promote biodiversity of native vegetation and wildlife through habitat restoration and protection.

The construction of the 876.32-acre Bank will be done in 4 phases due to size and logistics of grading and in-water work. Construction actions include: removing 100 years of farm infrastructure including fencing, gates, roads, duck blinds, and water pipes; levee and water control structure removal and modification for floodplain reconnection, tidal hydrology and fish passage restoration. Fill will be removed to restore tidal and distributary channel morphology, and ditches will be filled. Invasive reed canary grass and other non-native species will be removed, lowering floodplain elevations to increase inundation and promote native plant communities. Elevations will be modified to increase topographic diversity and support native woody and emergent plant communities. Aquatic habitat complexity will be increased through installation of large wood habitat structures. Oregon white oak habitats will be restored by removing competing tree species that are crowding the oak and competing for light and space, and new Oregon white oak habitat will be constructed to increase acreage of oak savannah and wet prairie.

To support permitting of the bank construction and updates to the Clark County Shorelines Master Plan this assessment documents the state and local shoreline jurisdiction of the Washington Department of Ecology (Ecology), Clark County (County) and the separate federal jurisdiction of the U.S. Army Corps of Engineers (Corps) as it relates to the Ordinary High Water Mark (OHWM) for Section 10 of the Rivers and Harbors Act. Wetlands and waterbodies within the Bank property are documented separately in a 2016 report by Cascade Environmental Group, titled "Plas Newydd Farm Wetlands and Other Waters Delineation Report, prepared for Plas Newydd LLC. Plas Newydd LLC received a letter of concurrence in the form of a jurisdictional determination (JD) from the U.S. Army Corps of Engineers (Corps) dated 6 September 2018. The delineation report and JD are provided under separate cover due to size.

The proposed 876-acre Bank is located wholly on privately owned property, Plas Newydd Farm which is owned by Plas Newydd LLC, in north Clark County, Washington (Figure 1). PN Farm and the Wapato Valley Bank are in Water Resource Inventory Area (WRIA) 27, the Lewis River watershed in the Columbia River basin, within the freshwater tidally influenced portion of the lower floodplain

at the confluence of the Lewis River at River Mile (RM) 87. The Bank is located approximately two-thirds of the distance between the mouth of the Columbia River as it enters the Pacific Ocean (RM 0) and Bonneville Dam (RM 146), which is the most downstream of 14 mainstem dams on the Columbia River. The Bank is situated west of U.S. Interstate 5 (I-5), east of the Columbia River, north of the town of Ridgefield, and south of the town of Woodland; in portions of Sections 1, 2, 11, and Donation Land Claim (DLC) 37<sup>1</sup>, and Section 12 in Township 4 North, Range 1 West (Clark County 2015; AINW, Inc. 2013). The situs address of PN Farm and Wapato Valley Bank is 33415 NW Lancaster Road, Ridgefield, Washington, 98642. The Bank encompasses 876.32 acres and is comprised of portions of Clark County tax parcel numbers 217593000, 217798000, and 218003000. The Bank is bordered by the BNSF Railway to the east, the Lewis River to the north, the Columbia River to the west, and Gee Creek and the Ridgefield National Wildlife Refuge (RNWR) to the south.

PN Farm is currently managed for sustainable family forestry, agriculture, and leased duck hunting. The land is topographically diverse and ranges in elevation from about 6 to 80 feet NAVD88. The site is hydrologically complex and influenced by the confluence setting, twice-daily backwater tidal influence from the Columbia River, seasonal flooding, and groundwater and hyporheic interactions. The Bank consists of diked and undiked wetlands (including open water lake, stream, and river channel; mudflat; emergent, low, and high marsh; wet pasture; scrub-shrub; and forested wetland), and uplands (including upland pasture, grassland, mixed deciduous/conifer forest, oak woodland, riparian forest, conifer forest, and dike/levee structure). The site supports biologically diverse habitats and native fish and wildlife species, including rare native plant communities and multiple special-status species.

## 2 METHODS

This assessment was prepared by Plas Newydd LLC staff. Kelley Jorgensen is the Plas Newydd President of Conservation and lead restoration ecologist responsible for the planning, development, and implementation of aquatic and terrestrial habitat restoration projects on 1000+ acres. She is leading the development and approval of the proposed 876-acre Wapato Valley Wetland Mitigation and Conservation Bank. With over 28 years of experience in the Pacific Northwest in applied ecology, Kelley's career to date has spanned the public, private and non-profit sectors. She combines her expertise in Pacific Northwest watershed ecology, field biology, interdisciplinary restoration approaches, environmental project management, permitting and facilitation to lead the Conservation Program in restoring this dynamic, complex and biodiverse landscape.

Chris Watson, a certified GISP, is Plas Newydd's GIS analyst, field geologist and data manager. His background includes over 20 years in the Pacific Northwest

---

<sup>1</sup> Sometimes shown as DLC 57, which varies by data source due to Donation Land Claim origin.



permitting and regulatory consulting environments. Chris provides the Conservation Program team with hydrologic and other modeling as well as GIS analytical capabilities. Chris is adept at bringing to bear the correct spatial data and analyses to solve complex and often multifaceted problems. He has a skillset that includes project management, GIS analysis, geologic evaluation and exploration, technical writing, public education support, litigation support, computer simulations and modeling, and database design. Mr. Watson has spent the last six years working on river and habitat restoration projects in the lower Columbia. Chris has been part of over 20 NEPA project teams in Oregon, Washington, Idaho, and Utah.

Sophie Ernst is a field biologist and is a Certified Erosion and Sediment Control Lead, and certified in ArcGIS, with 4 years of environmental data collection and analysis. She is skilled in Real-Time Kinematic (RTK) Global Positioning System (GPS) and other remote sensing data collection and analysis, biotic and abiotic field data collection and analysis, identification of flora and fauna, collection and interpretation of hydrologic data, and use of Python, Bad Elf and Excel. Sophie has a Bachelor of Arts in Environmental Studies from the University of Washington, and a Geographic Information System (GIS) Certificate from Portland Community College.

Hannah Mortensen is a field biologist, is GIS-certified and a licensed Unmanned Aerial Vehicle (UAV, or drone) pilot, with over 4 years of environmental data collection and analysis. She is skilled in Real-Time Kinematic (RTK) Global Positioning System (GPS) and other remote sensing data collection and analysis, 3D modeling, biotic and abiotic field data collection and analysis, identification of flora and fauna, collection and interpretation of hydrologic data, and use of Python, Bad Elf and Excel. Hannah has a Bachelor of Science in Ecology from The Evergreen State College, and a Geographic Information System (GIS) Certificate from Portland Community College.

Karen Adams is a senior wetland ecologist and monitoring lead. She has over 25 years of experience in monitoring the health and status of watershed conditions, specializing in wetlands and aquatic habitats. Her work has focused on developing monitoring plans and protocols, statistical analysis of environmental and experimental data, and reporting. Karen has earned degrees in Environmental Science, Wetlands Biology, and Ecology and Evolutionary Biology, investigating the effects of channel modification for flood management on forested wetlands, and the interactions between native and invasive wetland plant species. She has worked in and around Washington State's salmon bearing ecosystems for the last 10 years for the Washington State Department of Ecology, the Lower Columbia Fish Recovery Board and Plas Newydd LLC.

Documentation, field data collection and hydrologic assessment methods for the OHWM determination are based on from "Determining the Ordinary High Water Mark for Shoreline Management Act Compliance in Washington State" (Ecology 2016). Extensive office and field assessments have been conducted

(many are ongoing) over a period of 5+ years (2014-2019) collecting biotic and abiotic data to document pre-project conditions on the 876.32 acre Bank and portions of the roughly 800 acres of Plas Newydd property in forestry outside the Bank. The data provided here is a summary of relevant information helpful to understand the OHWM determination and includes a combination of field indicators and a hydrologic (stream and tidal) assessment conducted for the Lewis River using the stream methodology, field indicators for Lancaster Lake, and a combination for the Columbia River using the marine or tidal methodology of mean higher high water and more traditional fluvial or stream field indicators Lewis River and Gee Creek; both stream and tidal methods in combination are the most useful for delineating tidal fresh waters. The office assessment provided is focused on the hydrologic assessment, detailed in the next section. PN Conservation Program staff identified 9.2 miles (48,630 lineal feet) of shoreline areas along 4 waterbodies located on or adjacent to PN Farm for delineation of OHWM including the Columbia River, Lewis River, Gee Creek, and Lancaster Lake (Table 1, Figure 2). Additional shoreline areas are located along Allen Creek (aka Allen Canyon Creek) and Lake Rosannah that are within the property boundary, however those areas were not identified for delineation as there are no proposed construction projects that could affect them at this time, nor do they appear to require updates or changes in the current 2019/2020 Clark County Shoreline Masterplan update process.

**Table 1. Waterbodies and Shoreline Areas included in Delineation of OHWM**

<b>Waterbody</b>	<b>River Miles</b>	<b>Miles of Shoreline</b>	<b>Lineal Feet of Shoreline</b>
Columbia River	87 – 87.3	0.45	2,405
Lewis River	0 – 2.75	4.55	24,045
Gee Creek	0 – 2.4	2.71	14,327
Lancaster Lake	N/A	1.49	7,853
<b>Total</b>	<b>5.45</b>	<b>9.2</b>	<b>48,630</b>

### 3 FIELD ASSESSMENT

Field visits focused on OHWM data collection were made at multiple locations along the above mentioned shorelines for the purpose of recording field indicators (vegetation, scour lines, wrack lines, flattened vegetation, soil markers, etc.) on the following dates:

- 1/9/2018
- 1/12/2018
- 1/15/2018
- 7/11/2019
- 7/12/2019
- 7/15/2019
- 7/16/2019
- 11/18/2019
- 11/19/2019

- 11/20/2019
- 12/2/2019
- 12/3/2019
- 12/4/2019

Plas Newydd technical staff collected field indicator and topographic elevation data at over 95 points scattered along 9.2 miles of shoreline. Field data points were concentrated in locations where Wapato Valley Bank proposed construction would overlap or approach OHW areas or where field indicators were the most easily discerned. Attachment A includes the field data forms and an overview map showing the locations of the RTK GPS data collection. Species (Latin) names and common names for vegetation discussed here are presented in tabular form in Attachment B. Vegetation, scour lines, bank erosion/channel scour, flattened vegetation from “drainage patterns” (tidal surge or fluvial flows), top of bank, overbank deposits and wrack lines were evident in various locations. Elevations were taken of OHWM features and analysis found patterns indicative of fluvial and/or tidal hydrologic influence, described further in the hydrologic assessment discussion and conclusions. Due to the large size of the shoreline area being delineated, patterns were found during field indicator and elevation data analysis and averages were used to create the OHWM across long stretches of shoreline.

### **3.1 COLUMBIA RIVER OHWM**

Field indicators are ephemeral, dynamic and highly variable in this mainstem lower Columbia River location, influenced by complex hydrodynamics including heavily-managed flows and regulated spill of the Columbia River hydropower system, tidal influence and backwater effects, and confluence effects from the Lewis River (also hydromodified by 3 channels-spanning hydroelectric dams upstream) and the Willamette River and Multnomah channel which enter the Columbia just upstream and across from the PN Farm property. The Columbia River is influenced by snow-melt driven spring freshet flows fed by the Rocky and Cascade mountain ranges which create short term but extreme rises in water surface elevation, sometimes on the order of 15 feet or more of fluctuation during a water year. High water on the Columbia is not typically in winter (which is the average high water for most west Cascade streams and rivers) but instead occurs between April and June.

The PN Farm property along the Columbia River is a rare low-elevation intact tidal surge plain with active erosion and accretion patterns and sand-dominated sediment transport. The shoreline is affected by fluvial flood flows, tidal backwater/slack tide conditions, fetch, and erosive wave action driven by wakes generated from a wide variety of vessel types ranging from very large ocean-going vessels with a deep draft to smaller fishing, pleasure and speed craft (including jet skis) which travel much closer to the shore and generate waves at a much higher frequency. To further complicate matters, soils are very sandy along the Columbia, groundwater hydrology is largely hyporheic and

wetlands have a high degree of upland plants depending upon the microclimate. Combined these elements serve to create a lot of “noise” and variation in elevation in the identification of field indicators.

The Columbia River (Clark County, WA side) shoreline on the western edge of the PN Farm property between approximately RM 87 and 87.3 (and the contiguous open sandy shoreline of the Lewis River confluence area) was surveyed over multiple site visits between January 2018 and December 2019. Field indicators were identified readily during both winter and summer (both seasons with prolonged low water conditions and strong tidal signal) that represent the lower limit of the OHWM including toe of lowest terrace, drainage patterns as shown by flattened vegetation, aquatic plants, and aquatic animals. Lower limit indicators fell within about one vertical foot of each other and were easily averaged. Field indicators for the upper limits were more difficult to discern and varied greatly in elevation due to lack of fixed objects, a site with little topographic relief and heavy wave action from vessel wakes. Upper limit indicators varied by 3 vertical feet and were more difficult to average as a result. See the hydrologic assessment for a discussion of mean higher high water, a datum relevant for this tidally dominated setting. Table 2 lists the dominant species of vegetation identified and their distribution across the OHWM gradient. The list identifies the dominant species identifiable at the time of survey but is not exhaustive.

**Table 2. Plant Distribution across Columbia River OHWM Gradient**

<b>Below OHWM</b>	<b>At/Straddling OHWM</b>	<b>Above OHWM</b>
Needle Spikerush, OBL	Reed canarygrass, FACW	Oregon ash, FACW
Softstem Bulrush, OBL	Willow sp., FACW (colonizing)	Willow sp, FACW (mature)
Slough Sedge, OBL	False indigo bush, FAC	Black cottonwood, FAC
Woolgrass, OBL	Red-osier dogwood, FACW	Himalayan blackberry FAC
	Rough cocklebur, FAC	Black hawthorn, FAC

### 3.2 LEWIS RIVER OHWM

The south shore of the Lewis River between RM 0 and 2.75 along PN Farm property was surveyed at 40 data points in 6 locations between January 2018 and December 2019. Much of shoreline of the Lewis River in the lower 3 miles is dominated by a persistent erosion-resistant clay with naturally steep banks and overlays of intermittent sandy benches. Some shoreline armoring (native basalt – ballast to 1-man rock in size) is also present in patches along the toe of the Lewis River levee between RM 1 up to RM 2 where Allen Creek flows into the Lewis River through twin culverts. Field indicators identified include scour/moss line on rocks, sediment lines on rocks, lack of soil horizons, aquatic plants, aquatic animals, vegetation changes, stain lines on fixed objects, depositional sediment changes, well developed soil horizons, relic floodplain surface, exposed

roots/root scour, bank erosion, wrack lines and benches. Field indicators generally fell within 12-18 inches of each other and were logical when averaged across the 4.5 miles of shoreline surveyed.

**Table 3. Plant Distribution across Lewis River OHWM Gradient**

Below OHWM	At/Straddling OHWM	Above OHWM
Sedge sp, OBL	Reed canarygrass, FACW	Oregon ash, FACW
Rush sp, OBL	Red-osier dogwood, FACW	Oregon white oak, FACU/UPL
	Western goldenrod, FACW	Black cottonwood, FAC
		Himalayan blackberry, FAC

### 3.3 GEE CREEK OHWM

The north shore of Gee Creek between RM 0 and 2.4 along PN Farm property was surveyed at 24 data points in 4 locations between January 2018 and December 2019. The shoreline of Gee Creek is dominated by either a persistent erosion-resistant clay with naturally steep banks or naturally occurring native basalt outcrops. A narrow rock wall canyon also exists about halfway along the surveyed length. Field indicators identified include scour/moss line on rocks, sediment lines on rocks, lack of soil horizons, clean cobbles/boulders, aquatic plants, aquatic animals, vegetation changes, stain lines on fixed objects, depositional sediment changes, well developed soil horizons, relic floodplain surface, exposed roots/root scour, bank erosion, wrack lines and benches. Field indicators generally fell within 12-18 inches of each other and made sense when averaged across the 2.7 miles of shoreline surveyed.

**Table 4. Plant Distribution across Gee Creek OHWM Gradient**

Below OHWM	At/Straddling OHWM	Above OHWM
Sedges, OBL	Reed canarygrass, FACW	Oregon ash, FACW
Needle spikerush, OBL	Red-osier dogwood, FACW	Oregon white oak, FACU/UPL
Wapato, OBL	Western goldenrod, FACW	Black cottonwood, FAC
	Moss sp., UPL	Douglas-fir, FACU
	Stonecrop, UPL	Himalayan blackberry FAC
	Willow sp. FACW	Snowberry, FACU

### 3.4 LANCASTER LAKE OHWM

Lancaster Lake is a perennially ponded impounded area created by a channel spanning dike (the Narrows dike) that isolates a large historic floodplain area

from Gee Creek to the south, and the Lewis River to the north is separated by another levee system. The dike has one small tide gate with a flapper valve that prevents Gee Creek from backwatering into the floodplain and Lancaster Lake, but allows some discharge out of the lake through the tidegate when water surface elevations in Lancaster Lake are higher than Gee Creek. The lake is largely fed by hyporheic groundwater because it is in the Columbia and Lewis River floodplains, and from precipitation and seeps. Water level monitoring inside and outside the levee has demonstrated that Lancaster Lake generally tracks the water levels in the Columbia during spring freshet fluctuations and flood flows from floodplain recharge with delays in both runup and flood recession. The unique floodplain setting creates a challenging location to determine the upper limit of the OHWM towards the extensive associated wetlands within the broad flat floodplain to the north of the lake. The lake is bounded to the east and west by naturally occurring basalt outcrops and bounded to the south by the Narrows levee, which is also armored with native locally sourced basalt levee rock, that show more obvious field indicators for the upper limit of the OHWM.

Twenty-two data points were taken in 4 locations along 1.5 miles of Lancaster Lake shoreline between July and December 2019. Field indicators documented include vegetative changes, sediment deposits, clean cobbles/bedrock, lack of soil horizon, aquatic plants, aquatic animals, and water marks on the shoreline and downed large wood, and a review of time series imagery that captured annual highwater events. From the documented field indicators, the OHWM is a relatively vertically and horizontally wide zone that spans across a gradation of more than four feet between the upper and lower limits. The OHWM was averaged across the upper limit indicator elevations, which generally fell within 12 – 18 inches of each other. Table 5 lists the dominant species of vegetation and their distribution across the OHWM gradient. The list identifies the dominant species recorded at the time of survey but is not exhaustive. Attachment A includes a map of locations of the data points and field data forms.

**Table 5. Plant Distribution across Lancaster Lake OHWM Gradient**

<b>Below OHWM</b>	<b>At/Straddling OHWM</b>	<b>Above OHWM</b>
Wapato, OBL	Reed Canarygrass, FACW	Oregon White Oak, FACU
Polygonum Species, OBL	Salix Sp, FACW	Douglas-Fir, FACU
Reed Canarygrass, FACW	Douglas Spirea, FACW	Vine Maple, FAC
Bull Rush, OBL	Oregon Ash, FACW	Himalayan blackberry, FAC
Rough cocklebur, FAC	Herb Robert, FACU	Scot's Broom, NI
Sparganium sp., OBL	Birdsfoot trefoil, FACU	Licorice fern, NI
		Camas, FACW

## 4 HYDROLOGIC ASSESSMENT METHODS

This section summarizes the methods, data, and results used in hydrologic assessments of the Wapato Valley project and PN Farm shorelines areas. As the location has both stream (fluvial) and tidal freshwater shoreline areas, this report includes hydrologic assessments of each. The hydrologic assessments were performed in conjunction with and supplementary to OHWM field assessment of the same shorelines, described above.

Wapato Valley lies in the floodplain at the confluence of the Lewis River WRIA 27 with the mainstem Columbia River at RM 87. Wapato Valley is located in the freshwater tidal zone and experiences a daily tidal range of 2–4 feet on average (NOAA 2011). Due to the complexity of the hydrologic conditions at Wapato Valley, it cannot be classified as simply “high energy” or “low energy.” PN Farm includes 9.2 miles of shoreline (Wapato Valley includes subset of that) (Table 1). Lancaster Lake has no fluvial in-flow with shorelines mainly affected by a subdued reflection in water surface level of that in the Columbia River. Gee Creek has shorelines with both a backwater area that is open and punctuated with abrupt hard-rock islands and a constricted channel bounded by mostly erosion-resistant consolidated clay or bedrock shore. Flow in Gee Creek is in both directions up and downstream depending mainly on the Columbia River WSL and tides. The Columbia River shoreline within Wapato Valley transitions from an aggrading shore near the mouth of Gee Creek to an eroding shoreline at the mouth of the Lewis River. Shores on the Lewis River portion of Wapato Valley exhibit high energy erosion characteristics near the mouth with lower energy characteristics upstream.

WRIA 27 encompasses over 1,300 square miles and drains the western slope of the Cascade Mountain range, emptying into the Columbia River at river mile 87 (Corps 2014). Downstream flow on the Lewis River is regulated by the three upstream hydroelectric dams and reservoir systems, fish protection instream flow rules, and various water management strategies (Ecology 2016a).

The Columbia River is approximately 1,243 miles in length and drains over 258,000 square miles in seven states, and one Canadian province. Flow in the Columbia River is regulated by 14 major dams in the main stem and 46 in its tributaries (NRC 2004). Flows in the lower Columbia River are highly modified by the upstream water control structures, the geographic extent and complexity of its basin, water management practices, power generation, and other factors. Columbia River shorelines within Wapato Valley are directly affected by dynamically changing WSL and flows dictated by daily tides, commercial ship traffic, and upriver spill control facilitating power generation, agriculture needs, flood control, and fish migration. Fluctuations also occur from year to year based on snow pack, precipitation levels, and local climate changes.

## 5 STREAM HYDROLOGIC ASSESSMENT

The stream and tidal hydrology assessment methods provided by the Washington Department of Ecology in Publication no. 16-06-029 (Ecology 2016) analyze stream flow data from proximal or surrogate stream gages. The goal of these analyses is to provide context and to capture the flow range also referred to as “bookend” values. Context can be useful in spotting trends or events that may otherwise obscure the indicators in the field, as is the case along the shorelines of the rivers and streams within Wapato Valley. Conversely, analyzing the recent and historic flows can help in planning field efforts around a time when indicators are most likely to be found. The flow range or “bookend” data is useful in bracketing elevation ranges to inform on-site OHWM field assessments and cross-checking field-driven determination results.

### 5.1 STEP 1 AND 2: USE GAGE DATA TO APPROXIMATE UPPER AND LOWER EXTREMES FOR OHW FLOWS AND CORRELATE TO STAGE

The nearest gage on the Lewis River is USGS 14220500 located in Ariel, WA at 45.95194° N, 122.5628° W. The Ariel, WA gage is approximately 18 miles upstream from Wapato Valley and has been recording from July 1, 1909 until the present (USGS 2019) (Figure 3). The channel at the gage location is approximately 235 feet wide at a stage of 10 feet. The upstream dams were finalized in 1958; consequently, the analysis uses data from 1958 to present as it most accurately reflects current flow conditions.

#### 5.1.1 Generate the upper bookends by estimating the two-year peak and minimum peak flow

Using the downloaded dataset, the calculated median is 24,800 cubic feet per second (cfs) corresponding to a stage of 11.8 feet. The minimum peak flow is 9,670 cfs corresponding to a stage of 6.54 feet. The chart method results were cross-checked with the spreadsheet method and found to match (Figure 4 and Table 6).

**Table 6. Maximum peak annual discharge data 1958–2017 Lewis River (aka “spreadsheet method”).**

Date	cfs	Stage (ft)
1958-02-12	18,300	10.52
1959-01-24	32,800	15.12
1959-10-12	21,400	11.33
1960-11-24	48,200	19.3
1961-12-20	11,900	7.72
1962-11-20	75,500	25.7
1964-01-25	17,700	9.98
1964-12-22	44,000	17.49
1966-08-01	11,900	7.76
1966-12-13	50,500	19.12



Date	cfs	Stage (ft)
1968-02-23	31,100	14.02
1968-11-11	21,000	11.03
1970-01-23	41,800	16.96
1971-01-25	23,300	11.76
1972-03-13	36,400	15.55
1972-12-24	18,000	9.99
1974-01-15	59,600	21.13
1975-01-14	22,400	11.46
1975-12-04	64,500	22.63
1976-12-02	11,800	7.61
1977-12-02	71,900	24.38
1978-11-15	11,800	7.62
1980-01-12	12,000	7.71
1980-12-26	53,700	19.93
1982-02-20	40,700	16.67
1983-01-07	27,000	12.78
1983-11-17	17,100	9.5
1985-06-07	22,100	11.29
1986-02-24	27,700	13.06
1986-11-24	12,100	7.53
1987-12-10	12,300	7.61
1989-02-06	11,700	7.51
1990-01-10	42,000	16.85
1990-11-25	39,600	16.23
1992-01-30	12,600	7.68
1993-04-03	12,000	7.49
1994-01-08	11,800	7.45
1995-02-20	26,600	12.56
1996-02-08	86,400	27.38
1997-01-01	34,100	14.92
1997-11-21	12,200	7.63
1998-12-29	35,900	15.43
1999-12-15	35,700	15.37
2001-05-14	9,670	6.54
2001-12-17	14,700	8.6
2003-01-31	49,300	18.98
2004-01-29	11,700	7.44
2005-01-17	16,500	9.3
2006-01-11	29,900	13.68
2006-11-06	39,900	16.54

Date	cfs	Stage (ft)
2007-12-04	18,200	9.89
2009-01-07	40,300	16.63
2010-01-05	12,700	7.79
2011-01-16	35,400	15.22
2011-12-29	17,900	9.66
2012-11-20	22,900	11.4
2014-03-09	26,400	12.53
2014-11-27	16,700	9.25
2015-12-11	31,700	14.14
2017-03-16	26,300	12.48
<b>Peak High (median)</b>	<b>2,4800</b>	--
<b>Peak Low (minimum)</b>	<b>9,670</b>	--

### 5.1.2 Refine the Range

To refine the vertical range, the upper limit or “bookend” flow is reduced to a flow value that is exceeded at least once each year in 60 percent of years. A plot and table of the daily mean discharge and stage were pulled for 2002–2017 with 16 years represented. The calculated value using the iterative method in the spreadsheet was 16,400 cfs. A flow 16,400 cfs meets the criteria of being exceeded in 60% of the years in the analysis data set. The 16,400 cfs peak flow, which corresponds to a stage of 9.15 feet, was exceeded 10 out of the 16 years or 62.5% of the years in the analysis dataset (Table 3 and Figure 3).

**Table 7. Number of times 16,400 cfs was exceeded in each year 2002–2017.**

Year	Exceedance Count
2002	0
2003	3
2004	0
2005	0
2006	9
2007	1
2008	1
2009	4
2010	0
2011	349
2012	4
2013	0
2014	5
2015	12
2016	0
2017	6

The lower limit or “bookend” value was raised slightly to 10,900 cfs corresponding to a stage of 6.95 feet. This adjustment was made to reduce the number of long duration exceedance events of previous value. The correlation of discharge to stage was done in both the spreadsheet and graphically. A correlation of discharge and stage is shown in Figure 5.

### 5.1.3 Step 3: Compare recent events to OHWM bookends

To identify recent discharge or flow events that may have left fresh indicators on the Wapato Valley site, daily gage data for the last 12 months was reviewed. It was determined that the lower bookend value was exceeded twice in the last 12 months with a stage of approximately 7.75 feet (Figure 6).

### 5.1.4 Stream assessment conclusions

Given the location of the Wapato Valley at the confluence of the Columbia and Lewis rivers, the distance (18 miles) downstream from the Ariel, WA gage, and the dynamic and complex nature of the site, the hydrologic stream assessment in this case is useful only as context for upstream basin contributions, but is not indicative of the holistic picture of the hydrologic conditions or influences on shoreline OHW conditions. In addition, the Lewis River hydrology at the Wapato Valley location is dominated and obscured by flood flows and tidal backwater flows from the mainstem Columbia River. The stream assessment does however clearly give a couple of windows of time (December 19–21 and 30–31, 2018) that we can use to correlate with tidal station data from the tidal assessment to focus the field assessment on the most probable local elevations.

## 6 TIDAL HYDROLOGIC ASSESSMENT

This hydrologic assessment is intended to be used in conjunction with the stream hydrologic assessment above to inform the OHWM determination at Wapato Valley. The tidal hydrology assessment methods provided in Ecology (2016b) help focus the field assessment by providing a range of elevations on the ground where field indicators are most likely to be found. The OHWM in most cases is based on observable field indicators and is always above the mean higher high water (MHHW). Tidal information should not be the sole basis for an OHWM determination; however, in locations where field indicators are missing or cannot be found at certain times of year, tidal data (MHHW) may be the only option for establishing the OHWM reliably and consistently (Ecology 2016; RCW 90.58.030(2)(c)). The OHW delineation document is conspicuously missing guidance on the very large area of freshwater tidal influence on the lower Columbia River.

### 6.1 STEPS 1–3: LOCATE AN APPROPRIATE STATION AND IDENTIFY TIDAL DATUMS

Wapato Valley is located at RM 87 on the Columbia River. The St. Helens, OR tidal station, ID 9439201, is located at RM 86. For the purposes of this assessment, all elevations from the St. Helens station will be given in Columbia River Datum (CRD) which is 4.28 feet less than NAVD 88 at this location. The MHHW at the St. Helens station is reported as 5.28 feet, which equates to 9.56 feet NAVD 88. The vertical offset of Wapato Valley from the St. Helens station is +0.2 feet, giving Wapato Valley a MHHW elevation of 9.76 feet NAVD 88 (NOAA 2011) (Table 4).

**Table 8. Local Datum Comparisons to MHHW at St. Helens Tidal Station.**

CRD (ft)	NAVD 88 +4.28 (ft)	Wapato Valley Upriver Offset +0.2 (ft NAVD 88)
5.28	9.56	9.76

It should be noted that MHHW is calculated on tidal epochs. A tidal epoch is the specific 19-year period adopted by the National Ocean Service as the official time segment over which tide observations are taken and reduced to obtain mean values (e.g., mean lower low water, etc.) for tidal datums. The present National Tidal Datum Epoch (NTDE) is 1983 through 2001 and is actively considered for revision every 20–25 years. The MHHW listed above for Wapato Valley is based on an epoch that ended in 2001 (NOAA 2011).

In the stream assessment, periods of peak flow were identified that have a higher probability of corresponding with the formation of OHWM indicators. When the St. Helens station data is correlated with the peak flow periods (December 19–21 and 30–31, 2018) identified in the stream assessment, water surface elevations from the St. Helens station are shown to peak from 5.5–8.3 feet CRD (9.98–12.78 feet NAVD 88). These hybrid bookends prove useful in identifying the OHWM on the Columbia and Lewis River shorelines at the Wapato Valley location.

## 6.2 TIDAL ASSESSMENT CONCLUSIONS

Given the hybridized fluvial-tidal nature and complex riverine setting at the confluence of the Columbia and Lewis rivers, and the tidal epoch date range from which the published MHHW was derived, the tidal assessment places the bookends between 5.5–8.3 feet CRD (9.98–12.78 feet NAVD 88) on the Lewis River shoreline portions of the Wapato Valley and between 2.05–5.28 feet CRD (6.53–9.76 feet NAVD 88) on the Columbia River shoreline sections of the site. As noted previously, the tidal assessment is meant to guide and supplement the field indicators assessment of the OHWM determination.

**Table 9. Hydrologic assessment “bookend” OHWM elevation ranges.**

Shoreline Location	Probable Low (CRD)	Probable Low (NAVD 88)	Probable High (CRD)	Probable High (NAVD 88)
Lewis River	5.5	9.98	8.3	12.78
Columbia River	2.05	6.53	5.28	9.76

## 7 CONCLUSIONS

The OHWM determination for the following four waterbodies located on or adjacent to the Plas Newydd LLC property pertaining to Plas Newydd Farm and Wapato Valley Bank, based on the analysis documented in this report through field indicators and hydrologic assessment are as follows:

**Table 10. OHWM Results for Plas Newydd Farm/Wapato Valley in NAVD88**

Columbia River	Lewis River	Gee Creek	Lancaster Lake
9.76 (MHHW)	11.8	11.8	10.57

## 8 REFERENCES

Corps (U.S. Army Corps of Engineers). 2014. CAD data: Oregon\_North\_Channel\_RM\_NAD83\_USft\_2014.dxf. U.S. Army Engineer District, Portland, Operations Division.

Ecology (Washington State Department of Ecology). 2016a. Focus on water availability: Lewis River Watershed, WRIA 27, Publication no. 11-11-031. <https://fortress.wa.gov/ecy/publications/documents/1111031.pdf> [Accessed 26 November 2019]

Ecology. 2016b. Determining the ordinary high water mark for Shoreline Management Act compliance in Washington State. Publication no. 16-06-029. <https://fortress.wa.gov/ecy/publications/SummaryPages/1606029.html> [Accessed 26 November 2019]

NOAA (National Oceanic and Atmospheric Administration). 2011. Tides and Currents: Tidal Datums at Columbia River, St. Helens, OR. <https://tidesandcurrents.noaa.gov/datums.html?id=9439201> [Accessed 26 November 2019]


NRC (National Research Council). 2004. Managing the Columbia River: Instream flows, water withdrawals, and salmon survival. Chapter 3. Hydrology and Water Management. The National Academies Press, Washington, D.C. <https://doi.org/10.17226/10962> [Accessed 26 November 2019]

USGS [United States Geologic Survey]. 2019. National Water Information System: Stream gage data for USGS 14220500 Lewis River at Ariel, Washington. [https://waterdata.usgs.gov/nwis/inventory?agency\\_code=USGS&site\\_no=14220500](https://waterdata.usgs.gov/nwis/inventory?agency_code=USGS&site_no=14220500) [Accessed 26 November 2019]

---


## FIGURES

---




**Figure 1. Property Overview & Vicinity**

Plas Newydd, LLC  
Conservation Program




-  Property Boundary
-  Wapato Valley Mitigation and Conservation Bank
-  Dike or Levee
-  River Mile Marker



**PLAS NEWYDD FARM**  
EST. 1941  
CONSERVATION PROGRAM


**OHWM Determination**






**Figure 2. OHWM Elevations**

Wapato Valley  
Mitigation and Conservation Bank



**OHWM Elevation (NAVD 88)**

- 9.76 - Columbia River
- 11.8 - Lewis River
- 11.8 - Gee Creek
- 10.57 - Lancaster Lake



PLAS NEWYDD FARM  
EST. 1941  
CONSERVATION PROGRAM

**OHWM Determination**

Figure 3.  
Location of the nearest tidal station and stream gage to Plas Newydd Farm and Wapato Valley.




Figure 4.  
Hydrograph of the maximum peak annual discharge data for the Lewis River 2-year and 1.01-year peak flows depicted (aka "chart method").




Figure 5.  
 Stage for the determined flow range values plotted on aligned discharge and stage graphs.





Figure 6.  
Daily discharge plotted with refined OHWM bookend limits from refined analysis.



---

## ATTACHMENTS


---

---

**ATTACHMENT A**


**FIELD DATA FORMS AND MAPS**

---




### Ordinary High Water Locations

Wapato Valley  
Mitigation and Conservation Bank



● Field Data Points  
 Bank Area  
 LL - Lancaster Lake  
 GC - Gee Creek  
 LR - Lewis River  
 CR - Columbia River



PLAS NEWYDD FARM  
EST. 1941  
CONSERVATION PROGRAM



# Appendix A: Field data form

## General Information

Site/Project: Wapato Valley  
 Name/Owner: Plas Newydd Farm  
 Location: Columbia River  
 Description: 45.85184, -122.777552  
points: CR-A-(1-3)

The following field form is for use in the field to help in making ordinary high water mark delineations on streams. The form should be used as a guide. A team consisting of a hydrologist/ geomorphologist and a biologist may be needed to accurately determine the ordinary high water mark.

## General Observations: Day of Site Visit


Date of site visit:	20 NOV 2019		
Time of site visit:	11:10		
Weather conditions:	Full Sun		
Watershed development:	Highly developed <input checked="" type="radio"/>	Mod. Developed <input type="radio"/>	Undeveloped <input type="radio"/>
Reach development:	Highly developed <input checked="" type="radio"/>	Mod. Developed <input type="radio"/>	Undeveloped <input type="radio"/>
Recent site disturbance?	No <input checked="" type="radio"/>	Yes <input type="radio"/>	Describe:
Upstream flow control devices?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: Bunpilledam
Bank armoring at the site?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: opposite side on Oregon side
Bank armoring up or downstream?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: upstream
Observable tidal backwater?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	
In-water structures? (i.e. bridge pilings, railroad embankments)	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: pilings
Animals grazing in riparian zone?	No <input checked="" type="radio"/>	Yes <input type="radio"/>	Describe:
Observable beaver activity?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: Beaver chews

## Complete Vegetation Transects

- Use guidelines in Chapter 4 to complete vegetation transects.
- Determine upper and lower bounds of the OHWM from vegetation transects.
- After completing vegetation transects, look for more field indicators near the upper and lower bounds of the OHWM. Use the checklist as guidance.

## Sketch

If a simple site, sketch a cross-sectional diagram of the site below. Include location of the waterway and upper and lower bounds of the OHWM defined by the vegetation communities or other OHWM indicators. Page 3 of the data form can be used for more complex sketches



## Additional Indicators

Check the indicators that are observable at the site that provide rationale for establishing the OHWM at this location. The rationale should be described in detail in the report and should be supported with photographs taken during the site visit.

	Soil and geomorphic indicators <sup>24</sup>	Vegetative indicators <sup>25</sup>	Other indicators
<b>Below OHWM</b>	<ul style="list-style-type: none"> <li>○ Sediment bars</li> <li>○ Scour line</li> <li>○ Clean cobbles/boulders.</li> <li>○ Bank erosion/scour</li> <li>○ Lack of soil horizons</li> </ul>	Vegetation tolerant of inundation or high flow disturbances such as: <ul style="list-style-type: none"> <li>○ Willows</li> <li>○ Black cottonwood</li> <li>○ Japanese knotweed</li> <li>○ Skunk cabbage</li> <li>○ Aquatic plants</li> </ul>	<ul style="list-style-type: none"> <li>○ Exposed roots/root scour</li> <li>○ Drainage patterns, as shown by flattened vegetation</li> <li>○ Aquatic animals</li> <li>○ Algal mats</li> <li>○ Iron staining</li> </ul>


<sup>24</sup> Refer to Chapter 4 for a more complete description of indicators.

<sup>25</sup> Species are provided as examples. Refer to Appendix B for a more complete listing of plant species and their distribution across the OHWM gradient. Some species occur in more than one category depending on site conditions. For example Indian plum and red alder may straddle the OHWM where soil drainage is high. They may occur above OHWM where soil drainage is low to moderate.

	Soil and geomorphic indicators <sup>24</sup>	Vegetative indicators <sup>25</sup>	Other indicators
At or straddling OHWM	<ul style="list-style-type: none"> <li>○ Top of bank</li> <li>✗ Toe of lowest terrace (if terrace has developed horizons which may include a duff layer and A and B horizons versus freshly deposited alluvium)</li> <li>○ Benches</li> </ul>	<ul style="list-style-type: none"> <li>✗ Willows</li> <li>○ Western red cedar</li> <li>○ Vine maple (streams)</li> <li>○ Black cottonwood</li> <li>○ Red alder</li> <li>○ Salmonberry (red canopy)</li> <li>○ Nootka rose</li> <li>○ Maidenhair and lady fern</li> <li>○ Blackberries soft stem</li> <li>○ Dune grasses bull rush</li> </ul>	<ul style="list-style-type: none"> <li>○ Sediment lines on vegetation or other fixed objects</li> <li>○ Change from channel deposits to older alluvium.</li> <li>✗ Darker stain lines on fixed objects</li> <li>○ Exposed roots/root scour.</li> <li>✗ Drainage patterns, as evidenced by flattened vegetation</li> <li>✗ Weathered and buried driftwood</li> </ul>
Above OHWM	<ul style="list-style-type: none"> <li>○ Hillslope toe</li> <li>✗ Terraces or alluvium with an organic horizon or other developed soil horizons</li> <li>✗ Relic floodplain surface</li> <li>✗ Well developed soil A and B horizons/duff layer</li> </ul>	<ul style="list-style-type: none"> <li>○ Indian plum</li> <li>○ Red alder</li> <li>○ Western red cedar</li> <li>○ Douglas fir</li> <li>○ Western hemlock</li> <li>○ Ponderosa pine</li> <li>○ Oregon white oak</li> <li>○ Coast pine willow</li> <li>○ Quaking aspen</li> <li>○ Vine maple (lakes)</li> <li>○ Blackberries black cottonwood</li> </ul>	<ul style="list-style-type: none"> <li>○ Lighter or no staining on fixed objects</li> <li>○ Overbank deposits</li> </ul>

### Notes

The needle spike is flattened. The soft stem bull rush is the edge of the OHWM.



Plant Distribution Across OHWM Gradient					
Below <del>Above</del> OHWM		At/Straddling OHWM		Above OHWM	
needle spikerush	OBL	reed canarygrass	FACW	sand bar willow	FACW
wapato	OBL	willow sp.	FACW	Oregon ash	FACW
wool sedge	OBL			reed canarygrass	FACW
rice cutgrass	OBL				

20 Nov 2019  
Photo Direction: S  
WSEL: 8.51ft NAVD88

CR-A-(1-3)



← OHWM: Flattened vegetation

## Appendix A: Field data form

### General Information

Site/Project: Wapato Valley /  
 Name/Owner: Plus Newydd Farm  
 Location: Columbia River  
 Description: 45.850831, -122.7770629  
points: CR-B. (1-3)

The following field form is for use in the field to help in making ordinary high water mark delineations on streams. The form should be used as a guide. A team consisting of a hydrologist/ geomorphologist and a biologist may be needed to accurately determine the ordinary high water mark.

### General Observations: Day of Site Visit


Date of site visit:	11-20-2019		
Time of site visit:	11:30		
Weather conditions:	full sun		
Watershed development:	Highly developed <input type="radio"/>	Mod. Developed <input checked="" type="radio"/>	Undeveloped <input type="radio"/>
Reach development:	Highly developed <input checked="" type="radio"/>	Mod. Developed <input type="radio"/>	Undeveloped <input type="radio"/>
Recent site disturbance?	No <input checked="" type="radio"/>	Yes <input type="radio"/>	Describe:
Upstream flow control devices?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: Bonneville Dam
Bank armoring at the site?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: opposite shore
Bank armoring up or downstream?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe:
Observable tidal backwater?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	
In-water structures? (i.e. bridge pilings, railroad embankments)	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: pilings
Animals grazing in riparian zone?	No <input checked="" type="radio"/>	Yes <input type="radio"/>	Describe:
Observable beaver activity?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: fresh chewed sticks

### Complete Vegetation Transects

- o Use guidelines in Chapter 4 to complete vegetation transects.
- o Determine upper and lower bounds of the OHWM from vegetation transects.
- o After completing vegetation transects, look for more field indicators near the upper and lower bounds of the OHWM. Use the checklist as guidance.

## Sketch

If a simple site, sketch a cross-sectional diagram of the site below. Include location of the waterway and upper and lower bounds of the OHWM defined by the vegetation communities or other OHWM indicators. Page 3 of the data form can be used for more complex sketches



## Additional Indicators

Check the indicators that are observable at the site that provide rationale for establishing the OHWM at this location. The rationale should be described in detail in the report and should be supported with photographs taken during the site visit.

	Soil and geomorphic indicators <sup>24</sup>	Vegetative indicators <sup>25</sup>	Other indicators
<b>Below OHWM</b>	<ul style="list-style-type: none"> <li>○ Sediment bars</li> <li>○ Scour line</li> <li>○ Clean cobbles/boulders.</li> <li>○ Bank erosion/scour</li> <li>○ Lack of soil horizons</li> </ul>	Vegetation tolerant of inundation or high flow disturbances such as: <ul style="list-style-type: none"> <li>○ Willows</li> <li>○ Black cottonwood</li> <li>○ Japanese knotweed</li> <li>○ Skunk cabbage</li> <li>☑ Aquatic plants</li> </ul>	<ul style="list-style-type: none"> <li>○ Exposed roots/root scour</li> <li>☑ Drainage patterns, as shown by flattened vegetation.</li> <li>☑ Aquatic animals</li> <li>○ Algal mats</li> <li>○ Iron staining</li> </ul>

<sup>24</sup> Refer to Chapter 4 for a more complete description of indicators.


<sup>25</sup> Species are provided as examples. Refer to Appendix B for a more complete listing of plant species and their distribution across the OHWM gradient. Some species occur in more than one category depending on site conditions. For example Indian plum and red alder may straddle the OHWM where soil drainage is high. They may occur above OHWM where soil drainage is low to moderate.

	Soil and geomorphic indicators <sup>24</sup>	Vegetative indicators <sup>25</sup>	Other indicators
At or straddling OHWM	<ul style="list-style-type: none"> <li>○ Top of bank</li> <li>☒ Toe of lowest terrace (if terrace has developed horizons which may include a duff layer and A and B horizons versus freshly deposited alluvium)</li> <li>☒ Benches</li> </ul>	<ul style="list-style-type: none"> <li>○ Willows</li> <li>○ Western red cedar</li> <li>○ Vine maple (streams)</li> <li>○ Black cottonwood</li> <li>○ Red alder <i>Red canyon grass</i></li> <li>○ Salmonberry</li> <li>○ Nootka rose</li> <li>○ Maidenhair and lady fern</li> <li>○ Blackberries <i>Wood sedge</i></li> <li>○ Dunegrasses</li> </ul>	<ul style="list-style-type: none"> <li>☒ Sediment lines on vegetation or other fixed objects</li> <li>○ Change from channel deposits to older alluvium.</li> <li>☒ Darker stain lines on fixed objects</li> <li>○ Exposed roots/root scour.</li> <li>☒ Drainage patterns, as evidenced by flattened vegetation</li> <li>☒ Weathered and buried driftwood</li> </ul>
Above OHWM	<ul style="list-style-type: none"> <li>☒ Hillslope toe</li> <li>☒ Terraces or alluvium with an organic horizon or other developed soil horizons</li> <li>☒ Relic floodplain surface</li> <li>☒ Well developed soil A and B horizons/duff layer</li> </ul>	<ul style="list-style-type: none"> <li>○ Indian plum <i>willows</i></li> <li>○ Red alder <i>Oregon ash</i></li> <li>○ Western red cedar</li> <li>○ Douglas fir</li> <li>○ Western hemlock</li> <li>○ Ponderosa pine</li> <li>○ Oregon white oak</li> <li>○ Coast pine</li> <li>○ Quaking aspen</li> <li>○ Vine maple (lakes)</li> <li>○ Blackberries</li> </ul>	<ul style="list-style-type: none"> <li>☒ Lighter or no staining on fixed objects</li> <li>○ Overbank deposits</li> </ul>

Notes

The best OHWM is the wood sedge. The needle spike-rush is flatter and below the OHWM





Plant Distribution Across OHWM Gradient					
Below <del>Above</del> OHWM		At/Straddling OHWM		Above OHWM	
needle spikenush	OBL	wool sedge	OBL	sand bar willow	FACH
wapato	OBL	rice cutgrass	OBL	black cottonwood	NI
				reed canarygrass	PACH
				rough cocklebur	FAC

20 Nov 2019  
Photo Direction: N  
WSEL: 8.51ft NAVD88

CR-B-(1-3)



OHWL: Flattened vegetation →

# Appendix A: Field data form

## General Information

Site/Project: Wapato Valley /  
 Name/Owner: Plas Newydd Farm  
 Location: Columbia River  
 Description: 45.849933, -122.777538  
points: CR-D-(1-5)

The following field form is for use in the field to help in making ordinary high water mark delineations on streams. The form should be used as a guide. A team consisting of a hydrologist/ geomorphologist and a biologist may be needed to accurately determine the ordinary high water mark.

## General Observations: Day of Site Visit


Date of site visit:	12-4-2019		
Time of site visit:	12:25		
Weather conditions:	sun		
Watershed development:	Highly developed <input type="radio"/>	Mod. Developed <input checked="" type="checkbox"/>	Undeveloped <input type="radio"/>
Reach development:	Highly developed <input checked="" type="checkbox"/>	Mod. Developed <input type="radio"/>	Undeveloped <input type="radio"/>
Recent site disturbance?	No <input checked="" type="checkbox"/>	Yes <input type="radio"/>	Describe:
Upstream flow control devices?	No <input type="radio"/>	Yes <input checked="" type="checkbox"/>	Describe: Bonneville Dam
Bank armoring at the site?	No <input checked="" type="checkbox"/>	Yes <input type="radio"/>	Describe: opposite shore
Bank armoring up or downstream?	No <input type="radio"/>	Yes <input checked="" type="checkbox"/>	Describe:
Observable tidal backwater?	No <input type="radio"/>	Yes <input checked="" type="checkbox"/>	
In-water structures? (i.e. bridge pilings, railroad embankments)	No <input type="radio"/>	Yes <input checked="" type="checkbox"/>	Describe: pilings
Animals grazing in riparian zone?	No <input checked="" type="checkbox"/>	Yes <input type="radio"/>	Describe:
Observable beaver activity?	No <input checked="" type="checkbox"/>	Yes <input type="radio"/>	Describe:

## Complete Vegetation Transects

- Use guidelines in Chapter 4 to complete vegetation transects.
- Determine upper and lower bounds of the OHWM from vegetation transects.
- After completing vegetation transects, look for more field indicators near the upper and lower bounds of the OHWM. Use the checklist as guidance.

## Sketch

If a simple site, sketch a cross-sectional diagram of the site below. Include location of the waterway and upper and lower bounds of the OHWM defined by the vegetation communities or other OHWM indicators. Page 3 of the data form can be used for more complex sketches



## Additional Indicators

Check the indicators that are observable at the site that provide rationale for establishing the OHWM at this location. The rationale should be described in detail in the report and should be supported with photographs taken during the site visit.

	Soil and geomorphic indicators <sup>24</sup>	Vegetative indicators <sup>25</sup>	Other indicators
<b>Below OHWM</b>	<ul style="list-style-type: none"> <li>○ Sediment bars</li> <li>☒ Scour line</li> <li>○ Clean cobbles/boulders.</li> <li>☒ Bank erosion/scour</li> <li>☒ Lack of soil horizons</li> </ul>	Vegetation tolerant of inundation or high flow disturbances such as: <ul style="list-style-type: none"> <li>○ Willows</li> <li>○ Black cottonwood</li> <li>○ Japanese knotweed</li> <li>○ Skunk cabbage</li> <li>☒ Aquatic plants</li> </ul>	<ul style="list-style-type: none"> <li>☒ Exposed roots/root scour</li> <li>○ Drainage patterns, as shown by flattened vegetation.</li> <li>○ Aquatic animals</li> <li>○ Algal mats</li> <li>○ Iron staining</li> </ul>


<sup>24</sup> Refer to Chapter 4 for a more complete description of indicators.

<sup>25</sup> Species are provided as examples. Refer to Appendix B for a more complete listing of plant species and their distribution across the OHWM gradient. Some species occur in more than one category depending on site conditions. For example Indian plum and red alder may straddle the OHWM where soil drainage is high. They may occur above OHWM where soil drainage is low to moderate.

	Soil and geomorphic indicators <sup>24</sup>	Vegetative indicators <sup>25</sup>	Other indicators
At or straddling OHWM	<ul style="list-style-type: none"> <li>○ Top of bank</li> <li>○ Toe of lowest terrace (if terrace has developed horizons which may include a duff layer and A and B horizons versus freshly deposited alluvium)</li> <li>☒ Benches</li> </ul>	<ul style="list-style-type: none"> <li>○ Willows</li> <li>○ Western red cedar</li> <li>○ Vine maple (streams)</li> <li>○ Black cottonwood</li> <li>○ Red alder</li> <li>○ Salmonberry <i>MOF</i></li> <li>○ Nootka rose</li> <li>○ Maidenhair and lady fern</li> <li>○ Blackberries</li> <li>○ Dunegrasses</li> </ul>	<ul style="list-style-type: none"> <li>○ Sediment lines on vegetation or other fixed objects</li> <li>○ Change from channel deposits to older alluvium.</li> <li>☒ Darker stain lines on fixed objects</li> <li>☒ Exposed roots/root scour.</li> <li>○ Drainage patterns, as evidenced by flattened vegetation</li> <li>○ Weathered and buried driftwood</li> </ul>
Above OHWM	<ul style="list-style-type: none"> <li>○ Hillslope toe</li> <li>☒ Terraces or alluvium with an organic horizon or other developed soil horizons</li> <li>☒ Relic floodplain surface</li> <li>☒ Well developed soil A and B horizons/duff layer</li> </ul>	<ul style="list-style-type: none"> <li>○ Indian plum <i>dogwood</i></li> <li>○ Red alder</li> <li>○ Western red cedar</li> <li>○ Douglas fir</li> <li>○ Western hemlock <i>willow</i></li> <li>○ Ponderosa pine</li> <li>○ Oregon white oak</li> <li>○ Coast pine</li> <li>○ Quaking aspen</li> <li>○ Vine maple (lakes)</li> <li>☒ Blackberries</li> </ul>	<ul style="list-style-type: none"> <li>☒ Lighter or no staining on fixed objects</li> <li>○ Overbank deposits</li> </ul>

Notes

The best indicator of the OHWM at this location is the extent that moss can grow on the steeply eroded riverbanks.




Plant Distribution Across OHWM Gradient					
Below Above OHWM		At/Straddling OHWM		Above OHWM	
reed canarygrass	FACW	moss sp.	NI	Pacific willow	FACW
needle spikerush	OBL	himalaya blackberry	FACU	red-osier dogwood	FACW
				reed canarygrass	FACW
				himalaya blackberry	FACU
				black hawthorne	FAC

4 Dec 2019

Photo Direction: E

WSEL: 8.61ft NAVD88

CR-D-(1-5)



OHWM: Moss on upper limit

## Appendix A: Field data form

### General Information

Site/Project: Wapato Valley /  
 Name/Owner: PLAS NEWYDD FARM  
 Location: Gee Creek  
 Description: 45.846468, -122.75082  
points: GC-A-(1-9)

The following field form is for use in the field to help in making ordinary high water mark delineations on streams. The form should be used as a guide. A team consisting of a hydrologist/ geomorphologist and a biologist may be needed to accurately determine the ordinary high water mark.

### General Observations: Day of Site Visit

Date of site visit:	11-18-2019		
Time of site visit:	15:50		
Weather conditions:	light rain		
Watershed development:	Highly developed <input checked="" type="checkbox"/>	Mod. Developed <input type="checkbox"/>	Undeveloped <input type="checkbox"/>
Reach development:	Highly developed <input type="checkbox"/>	Mod. Developed <input checked="" type="checkbox"/>	Undeveloped <input type="checkbox"/>
Recent site disturbance?	No <input type="checkbox"/>	Yes <input type="checkbox"/>	Describe:
Upstream flow control devices?	No <input type="checkbox"/>	Yes <input checked="" type="checkbox"/>	Describe: Gee Creek bridge construction
Bank armoring at the site?	No <input type="checkbox"/>	Yes <input checked="" type="checkbox"/>	Describe: Basalt bluffs act as natural armoring.
Bank armoring up or downstream?	No <input type="checkbox"/>	Yes <input checked="" type="checkbox"/>	Describe: Gee Creek upstream of site.
Observable tidal backwater?	No <input type="checkbox"/>	Yes <input checked="" type="checkbox"/>	
In-water structures? (i.e. bridge pilings, railroad embankments)	No <input type="checkbox"/>	Yes <input checked="" type="checkbox"/>	Describe: Levee w/ tidegate to north block tide to Lancaster Lake.
Animals grazing in riparian zone?	No <input checked="" type="checkbox"/>	Yes <input type="checkbox"/>	Describe:
Observable beaver activity?	No <input type="checkbox"/>	Yes <input checked="" type="checkbox"/>	Describe: Channels and lodges.


### Complete Vegetation Transects

- Use guidelines in Chapter 4 to complete vegetation transects.
- Determine upper and lower bounds of the OHWM from vegetation transects.
- After completing vegetation transects, look for more field indicators near the upper and lower bounds of the OHWM. Use the checklist as guidance.



## Sketch

If a simple site, sketch a cross-sectional diagram of the site below. Include location of the waterway and upper and lower bounds of the OHWM defined by the vegetation communities or other OHWM indicators. Page 3 of the data form can be used for more complex sketches



## Additional Indicators

Check the indicators that are observable at the site that provide rationale for establishing the OHWM at this location. The rationale should be described in detail in the report and should be supported with photographs taken during the site visit.

	Soil and geomorphic indicators <sup>24</sup>	Vegetative indicators <sup>25</sup>	Other indicators
<b>Below OHWM</b>	<ul style="list-style-type: none"> <li>○ Sediment bars</li> <li>○ Scour line</li> <li>☒ Clean cobbles/boulders.</li> <li>○ Bank erosion/scour</li> <li>☒ Lack of soil horizons</li> </ul>	Vegetation tolerant of inundation or high flow disturbances such as: <ul style="list-style-type: none"> <li>○ Willows</li> <li>○ Black cottonwood</li> <li>○ Japanese knotweed</li> <li>○ Skunk cabbage</li> <li>☒ Aquatic plants</li> </ul>	<ul style="list-style-type: none"> <li>○ Exposed roots/root scour</li> <li>○ Drainage patterns, as shown by flattened vegetation.</li> <li>☒ Aquatic animals</li> <li>○ Algal mats</li> <li>○ Iron staining</li> </ul>

<sup>24</sup> Refer to Chapter 4 for a more complete description of indicators.


<sup>25</sup> Species are provided as examples. Refer to Appendix B for a more complete listing of plant species and their distribution across the OHWM gradient. Some species occur in more than one category depending on site conditions. For example Indian plum and red alder may straddle the OHWM where soil drainage is high. They may occur above OHWM where soil drainage is low to moderate.

	Soil and geomorphic indicators <sup>24</sup>	Vegetative indicators <sup>25</sup>	Other indicators
At or straddling OHWM	<ul style="list-style-type: none"> <li>○ Top of bank</li> <li>○ Toe of lowest terrace (if terrace has developed horizons which may include a duff layer and A and B horizons versus freshly deposited alluvium)</li> <li>○ Benches</li> </ul>	<ul style="list-style-type: none"> <li>○ Willows</li> <li>○ Western red cedar</li> <li>○ Vine maple (streams)</li> <li>○ Black cottonwood</li> <li>○ Red alder <i>break in</i></li> <li>○ Salmonberry <i>moss/stonecrop</i></li> <li>○ Nootka rose <i>to bank rock on</i></li> <li>○ Maidenhair and lady fern <i>bluffs</i></li> <li>○ Blackberries</li> <li>○ Dunegrasses</li> </ul>	<ul style="list-style-type: none"> <li>☒ Sediment lines on vegetation or other fixed objects</li> <li>○ Change from channel deposits to older alluvium.</li> <li>☒ Darker stain lines on fixed objects</li> <li>○ Exposed roots/root scour.</li> <li>○ Drainage patterns, as evidenced by flattened vegetation</li> <li>○ Weathered and buried driftwood</li> </ul>
Above OHWM	<ul style="list-style-type: none"> <li>○ Hillslope toe</li> <li>○ Terraces or alluvium with an organic horizon or other developed soil horizons</li> <li>○ Relic floodplain surface</li> <li>☒ Well developed soil A and B horizons/duff layer</li> </ul>	<ul style="list-style-type: none"> <li>○ Indian plum <i>Oregon ash</i></li> <li>○ Red alder</li> <li>○ Western red cedar</li> <li>☒ Douglas fir</li> <li>○ Western hemlock</li> <li>○ Ponderosa pine</li> <li>☒ Oregon white oak</li> <li>○ Coast pine</li> <li>○ Quaking aspen</li> <li>○ Vine maple (lakes)</li> <li>☒ Blackberries</li> </ul>	<ul style="list-style-type: none"> <li>○ Lighter or no staining on fixed objects</li> <li>○ Overbank deposits</li> </ul>

### Notes

The basalt rocks below the OHWM have very little to no vegetation on them

GC-A



Plant Distribution Across OHWM Gradient					
Below <del>Above</del> OHWM		At/Straddling OHWM		Above OHWM	
coon's tail	OBL	kane basalt		moss sp.	not listed
Eurasian watermilfoil	OBL	reed canarygrass	FACW	wormleaf stonecrop	not listed
				reed canarygrass	FACW
				hairy cat's ear	FACW
				cheatgrass	not listed
				camas	FACW
				Oregon ash	FACW
				Oregon white oak	FACW

18 Nov 2019

Photo Direction: N

WSEL: 7.01ft NAVD88

GC-A-(1-9)



← OHWM: Break in stonecrop/moss

# Appendix A: Field data form

## General Information

Site/Project: Wapato Valley /  
 Name/Owner: Plas Newydd Farm  
 Location: Gee Creek  
 Description: 45.24549, -122.751942  
points: GC-B - (1-7)

The following field form is for use in the field to help in making ordinary high water mark delineations on streams. The form should be used as a guide. A team consisting of a hydrologist/ geomorphologist and a biologist may be needed to accurately determine the ordinary high water mark.

## General Observations: Day of Site Visit


Date of site visit:	<u>11-19-2019</u>		
Time of site visit:	<u>10:00</u>		
Weather conditions:	<u>overcast</u>		
Watershed development:	Highly developed <input checked="" type="radio"/>	Mod. Developed <input type="radio"/>	Undeveloped <input type="radio"/>
Reach development:	Highly developed <input type="radio"/>	Mod. Developed <input checked="" type="radio"/>	Undeveloped <input type="radio"/>
Recent site disturbance?	No <input checked="" type="radio"/>	Yes <input type="radio"/>	Describe:
Upstream flow control devices?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: <u>Gee Creek bridge construction</u>
Bank armoring at the site?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: <u>Basalt bluffs act as natural armoring.</u>
Bank armoring up or downstream?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: <u>Gee Creek upstream of site</u>
Observable tidal backwater?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	
In-water structures? (i.e. bridge pilings, railroad embankments)	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: <u>Level w/ tidegate to north blocks tide to Lancaster Lake.</u>
Animals grazing in riparian zone?	No <input checked="" type="radio"/>	Yes <input type="radio"/>	Describe:
Observable beaver activity?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: <u>Channels and lodges.</u>

## Complete Vegetation Transects

- Use guidelines in Chapter 4 to complete vegetation transects.
- Determine upper and lower bounds of the OHWM from vegetation transects.
- After completing vegetation transects, look for more field indicators near the upper and lower bounds of the OHWM. Use the checklist as guidance.

## Sketch

If a simple site, sketch a cross-sectional diagram of the site below. Include location of the waterway and upper and lower bounds of the OHWM defined by the vegetation communities or other OHWM indicators. Page 3 of the data form can be used for more complex sketches



## Additional Indicators

Check the indicators that are observable at the site that provide rationale for establishing the OHWM at this location. The rationale should be described in detail in the report and should be supported with photographs taken during the site visit.

	Soil and geomorphic indicators <sup>24</sup>	Vegetative indicators <sup>25</sup>	Other indicators
<b>Below OHWM</b>	<ul style="list-style-type: none"> <li><input type="radio"/> Sediment bars</li> <li><input type="radio"/> Scour line</li> <li><input checked="" type="checkbox"/> Clean cobbles/boulders.</li> <li><input type="radio"/> Bank erosion/scour</li> <li><input checked="" type="checkbox"/> Lack of soil horizons</li> </ul>	Vegetation tolerant of inundation or high flow disturbances such as: <ul style="list-style-type: none"> <li><input type="radio"/> Willows</li> <li><input type="radio"/> Black cottonwood</li> <li><input type="radio"/> Japanese knotweed</li> <li><input type="radio"/> Skunk cabbage</li> <li><input checked="" type="checkbox"/> Aquatic plants</li> </ul>	<ul style="list-style-type: none"> <li><input type="radio"/> Exposed roots/root scour</li> <li><input type="radio"/> Drainage patterns, as shown by flattened vegetation.</li> <li><input checked="" type="checkbox"/> Aquatic animals</li> <li><input checked="" type="checkbox"/> Algal mats</li> <li><input type="radio"/> Iron staining</li> </ul>


<sup>24</sup> Refer to Chapter 4 for a more complete description of indicators.

<sup>25</sup> Species are provided as examples. Refer to Appendix B for a more complete listing of plant species and their distribution across the OHWM gradient. Some species occur in more than one category depending on site conditions. For example Indian plum and red alder may straddle the OHWM where soil drainage is high. They may occur above OHWM where soil drainage is low to moderate.

	Soil and geomorphic indicators <sup>24</sup>	Vegetative indicators <sup>25</sup>	Other indicators
At or straddling OHWM	<ul style="list-style-type: none"> <li>○ Top of bank</li> <li>○ Toe of lowest terrace (if terrace has developed horizons which may include a duff layer and A and B horizons versus freshly deposited alluvium)</li> <li>○ Benches</li> </ul>	<ul style="list-style-type: none"> <li>○ Willows</li> <li>○ Western red cedar</li> <li>○ Vine maple (streams)</li> <li>○ Black cottonwood</li> <li>○ Red alder</li> <li>○ Salmonberry</li> <li>○ Nootka rose</li> <li>○ Maidenhair and lady fern</li> <li>○ Blackberries</li> <li>○ Dunegrasses</li> </ul>	<ul style="list-style-type: none"> <li>☒ Sediment lines on vegetation or other fixed objects</li> <li>○ Change from channel deposits to older alluvium.</li> <li>☒ Darker stain lines on fixed objects</li> <li>○ Exposed roots/root scour.</li> <li>○ Drainage patterns, as evidenced by flattened vegetation</li> <li>○ Weathered and buried driftwood</li> </ul>
Above OHWM	<ul style="list-style-type: none"> <li>○ Hillslope toe</li> <li>☒ Terraces or alluvium with an organic horizon or other developed soil horizons</li> <li>☒ Relic floodplain surface</li> <li>☒ Well developed soil A and B horizons/duff layer</li> </ul>	<ul style="list-style-type: none"> <li>○ Indian plum <i>&amp; Oregon ash</i></li> <li>○ Red alder</li> <li>○ Western red cedar</li> <li>○ Douglas fir</li> <li>○ Western hemlock</li> <li>○ Ponderosa pine</li> <li>○ Oregon white oak</li> <li>○ Coast pine</li> <li>○ Quaking aspen</li> <li>○ Vine maple (lakes)</li> <li>☒ Blackberries</li> </ul>	<ul style="list-style-type: none"> <li>○ Lighter or no staining on fixed objects</li> <li>○ Overbank deposits</li> </ul>

## Notes

The high water line is visible on basalt bluffs where there is a break in moss growth. The nearby backwater area of Gee Creek is slightly higher and has filled in with sediment. It contains Oregon ash trees and a thick herb layer of reed canarygrass and native sedges. The top of the basalt bluff has very little soil and contains patches of Himalaya blackberry and snowberry. Upstream of the project site a construction project has Gee Creek dewatered for culvert replacement.



Plant Distribution Across OHWM Gradient						
Below						
<del>Above</del> OHWM		At/Straddling OHWM		Above OHWM		
wapato	OBL	reed	canarygrass	FACW	Oregon ash	FACW
needle spikerush	OBL				snowberry	FACU
hardstem bulrush	NI				himalaya blackberry	FACU
slough sedge	OBL					



19 Nov 2019

Photo Direction: W

WSEL: 8.87ft NAVD88

GC-B-(1-7)

OHWM: Vegetation change from wapato and sedge to upland vegetation



# Appendix A: Field data form

## General Information

Site/Project: Wapato Valley  
 Name/Owner: PLAS Newydd Farm  
 Location: Gee Creek  
 Description: 45.845924, -122.759277  
points: G6-C-(1-2)

The following field form is for use in the field to help in making ordinary high water mark delineations on streams. The form should be used as a guide. A team consisting of a hydrologist/ geomorphologist and a biologist may be needed to accurately determine the ordinary high water mark.

## General Observations: Day of Site Visit


Date of site visit:	20 Nov 2019		
Time of site visit:	14:50		
Weather conditions:	Full Sun		
Watershed development:	Highly developed <input type="radio"/>	Mod. Developed <input checked="" type="radio"/>	Undeveloped <input type="radio"/>
Reach development:	Highly developed <input type="radio"/>	Mod. Developed <input checked="" type="radio"/>	Undeveloped <input type="radio"/>
Recent site disturbance?	No <input checked="" type="radio"/>	Yes <input type="radio"/>	Describe:
Upstream flow control devices?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: culverts
Bank armoring at the site?	No <input checked="" type="radio"/>	Yes <input type="radio"/>	Describe:
Bank armoring up or downstream?	No <input checked="" type="radio"/>	Yes <input type="radio"/>	Describe:
Observable tidal backwater?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	
In-water structures? (i.e. bridge pilings, railroad embankments)	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe:
Animals grazing in riparian zone?	No <input checked="" type="radio"/>	Yes <input type="radio"/>	Describe:
Observable beaver activity?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: Beaver Chews

## Complete Vegetation Transects

- Use guidelines in Chapter 4 to complete vegetation transects.
- Determine upper and lower bounds of the OHWM from vegetation transects.
- After completing vegetation transects, look for more field indicators near the upper and lower bounds of the OHWM. Use the checklist as guidance.

## Sketch

If a simple site, sketch a cross-sectional diagram of the site below. Include location of the waterway and upper and lower bounds of the OHWM defined by the vegetation communities or other OHWM indicators. Page 3 of the data form can be used for more complex sketches



## Additional Indicators

Check the indicators that are observable at the site that provide rationale for establishing the OHWM at this location. The rationale should be described in detail in the report and should be supported with photographs taken during the site visit.

	Soil and geomorphic indicators <sup>24</sup>	Vegetative indicators <sup>25</sup>	Other indicators
<b>Below OHWM</b>	<ul style="list-style-type: none"> <li>○ Sediment bars</li> <li>○ Scour line</li> <li>○ Clean cobbles/boulders.</li> <li>○ Bank erosion/scour</li> <li><input checked="" type="checkbox"/> Lack of soil horizons</li> </ul>	Vegetation tolerant of inundation or high flow disturbances such as: <ul style="list-style-type: none"> <li>○ Willows</li> <li>○ Black cottonwood</li> <li>○ Japanese knotweed</li> <li>○ Skunk cabbage</li> <li><input checked="" type="checkbox"/> Aquatic plants</li> </ul>	<ul style="list-style-type: none"> <li>○ Exposed roots/root scour</li> <li><input checked="" type="checkbox"/> Drainage patterns, as shown by flattened vegetation.</li> <li><input checked="" type="checkbox"/> Aquatic animals</li> <li>○ Algal mats</li> <li>○ Iron staining</li> </ul>

reed canarygrass


<sup>24</sup> Refer to Chapter 4 for a more complete description of indicators.

<sup>25</sup> Species are provided as examples. Refer to Appendix B for a more complete listing of plant species and their distribution across the OHWM gradient. Some species occur in more than one category depending on site conditions. For example Indian plum and red alder may straddle the OHWM where soil drainage is high. They may occur above OHWM where soil drainage is low to moderate.

	Soil and geomorphic indicators <sup>24</sup>	Vegetative indicators <sup>25</sup>	Other indicators
At or straddling OHWM	<ul style="list-style-type: none"> <li><input checked="" type="checkbox"/> Top of bank</li> <li><input checked="" type="checkbox"/> Toe of lowest terrace (if terrace has developed horizons which may include a duff layer and A and B horizons versus freshly deposited alluvium)</li> <li><input type="checkbox"/> Benches</li> </ul>	<ul style="list-style-type: none"> <li><input type="checkbox"/> Willows</li> <li><input type="checkbox"/> Western red cedar</li> <li><input type="checkbox"/> Vine maple (streams)</li> <li><input type="checkbox"/> Black cottonwood</li> <li><input type="checkbox"/> Red alder</li> <li><input type="checkbox"/> Salmonberry <i>reel</i></li> <li><input type="checkbox"/> Nootka rose <i>canary</i></li> <li><input type="checkbox"/> Maidenhair and lady fern</li> <li><input type="checkbox"/> Blackberries <i>oregon</i></li> <li><input type="checkbox"/> Dunegrasses <i>ash</i></li> </ul>	<ul style="list-style-type: none"> <li><input checked="" type="checkbox"/> Sediment lines on vegetation or other fixed objects</li> <li><input checked="" type="checkbox"/> Change from channel deposits to older alluvium.</li> <li><input type="checkbox"/> Darker stain lines on fixed objects</li> <li><input type="checkbox"/> Exposed roots/root scour.</li> <li><input type="checkbox"/> Drainage patterns, as evidenced by flattened vegetation</li> <li><input type="checkbox"/> Weathered and buried driftwood</li> </ul>
Above OHWM	<ul style="list-style-type: none"> <li><input type="checkbox"/> Hillslope toe</li> <li><input checked="" type="checkbox"/> Terraces or alluvium with an organic horizon or other developed soil horizons</li> <li><input type="checkbox"/> Relic floodplain surface</li> <li><input checked="" type="checkbox"/> Well developed soil A and B horizons/duff layer</li> </ul>	<ul style="list-style-type: none"> <li><input type="checkbox"/> Indian plum</li> <li><input type="checkbox"/> Red alder</li> <li><input type="checkbox"/> Western red cedar</li> <li><input type="checkbox"/> Douglas fir</li> <li><input type="checkbox"/> Western hemlock</li> <li><input type="checkbox"/> Ponderosa pine</li> <li><input checked="" type="checkbox"/> Oregon white oak</li> <li><input type="checkbox"/> Coast pine <i>oregon</i></li> <li><input type="checkbox"/> Quaking aspen <i>ash</i></li> <li><input type="checkbox"/> Vine maple (lakes)</li> <li><input type="checkbox"/> Blackberries</li> </ul>	<ul style="list-style-type: none"> <li><input type="checkbox"/> Lighter or no staining on fixed objects</li> <li><input type="checkbox"/> Overbank deposits</li> </ul>

### Notes

Rocks on the edge of creek have a sediment line at the OHWM



Plant Distribution Across OHWM Gradient					
Below <del>Above</del> OHWM		At/Straddling OHWM		Above OHWM	
wapato	OBL	reed canarygrass	PACW	Oregon white oak	FACU
hardstem bulrush	NI			Oregon ash	FACW
reed canarygrass	FACW			snowberry	FACU

20 Nov 2019  
Photo Direction: E  
WSEL: 7.49ft NAVD88

GC-C-(1-2)



← OHWM: Sediment line on rocks

# Appendix A: Field data form

## General Information

Site/Project: Wapato Valley  
 Name/Owner: PLAS Newyald Farm  
 Location: Gee Creek  
 Description: 45.8411163, -122.760009  
points: GC-D-(1-3)

The following field form is for use in the field to help in making ordinary high water mark delineations on streams. The form should be used as a guide. A team consisting of a hydrologist/ geomorphologist and a biologist may be needed to accurately determine the ordinary high water mark.

## General Observations: Day of Site Visit


Date of site visit:	20 NOV 2019		
Time of site visit:	14:37		
Weather conditions:	Full Sun		
Watershed development:	Highly developed <input type="radio"/>	Mod. Developed <input checked="" type="radio"/>	Undeveloped <input type="radio"/>
Reach development:	Highly developed <input type="radio"/>	Mod. Developed <input checked="" type="radio"/>	Undeveloped <input type="radio"/>
Recent site disturbance?	No <input checked="" type="radio"/>	Yes <input type="radio"/>	Describe:
Upstream flow control devices?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: culverts
Bank armoring at the site?	No <input checked="" type="radio"/>	Yes <input type="radio"/>	Describe:
Bank armoring up or downstream?	No <input checked="" type="radio"/>	Yes <input type="radio"/>	Describe:
Observable tidal backwater?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	
In-water structures? (i.e. bridge pilings, railroad embankments)	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: Pilings
Animals grazing in riparian zone?	No <input checked="" type="radio"/>	Yes <input type="radio"/>	Describe:
Observable beaver activity?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: Beaver chews

## Complete Vegetation Transects

- Use guidelines in Chapter 4 to complete vegetation transects.
- Determine upper and lower bounds of the OHWM from vegetation transects.
- After completing vegetation transects, look for more field indicators near the upper and lower bounds of the OHWM. Use the checklist as guidance.

## Sketch

If a simple site, sketch a cross-sectional diagram of the site below. Include location of the waterway and upper and lower bounds of the OHWM defined by the vegetation communities or other OHWM indicators. Page 3 of the data form can be used for more complex sketches



## Additional Indicators

Check the indicators that are observable at the site that provide rationale for establishing the OHWM at this location. The rationale should be described in detail in the report and should be supported with photographs taken during the site visit.

	Soil and geomorphic indicators <sup>24</sup>	Vegetative indicators <sup>25</sup>	Other indicators
<b>Below OHWM</b>	<ul style="list-style-type: none"> <li><input type="checkbox"/> Sediment bars</li> <li><input type="checkbox"/> Scour line</li> <li><input checked="" type="checkbox"/> Clean cobbles/boulders.</li> <li><input type="checkbox"/> Bank erosion/scour</li> <li><input checked="" type="checkbox"/> Lack of soil horizons</li> </ul>	Vegetation tolerant of inundation or high flow disturbances such as: <ul style="list-style-type: none"> <li><input type="checkbox"/> Willows</li> <li><input type="checkbox"/> Black cottonwood</li> <li><input type="checkbox"/> Japanese knotweed</li> <li><input type="checkbox"/> Skunk cabbage</li> <li><input checked="" type="checkbox"/> Aquatic plants</li> </ul>	<ul style="list-style-type: none"> <li><input type="checkbox"/> Exposed roots/root scour</li> <li><input type="checkbox"/> Drainage patterns, as shown by flattened vegetation.</li> <li><input checked="" type="checkbox"/> Aquatic animals</li> <li><input type="checkbox"/> Algal mats</li> <li><input type="checkbox"/> Iron staining</li> </ul>

reed canary grass  
needle spikenish


<sup>24</sup> Refer to Chapter 4 for a more complete description of indicators.  
<sup>25</sup> Species are provided as examples. Refer to Appendix B for a more complete listing of plant species and their distribution across the OHWM gradient. Some species occur in more than one category depending on site conditions. For example Indian plum and red alder may straddle the OHWM where soil drainage is high. They may occur above OHWM where soil drainage is low to moderate.



	Soil and geomorphic indicators <sup>24</sup>	Vegetative indicators <sup>25</sup>	Other indicators
At or straddling OHWM	<ul style="list-style-type: none"> <li><input checked="" type="checkbox"/> Top of bank</li> <li><input checked="" type="checkbox"/> Toe of lowest terrace (if terrace has developed horizons which may include a duff layer and A and B horizons versus freshly deposited alluvium)</li> <li><input checked="" type="checkbox"/> Benches</li> </ul>	<ul style="list-style-type: none"> <li><input type="checkbox"/> Willows</li> <li><input type="checkbox"/> Western red cedar</li> <li><input type="checkbox"/> Vine maple (streams)</li> <li><input type="checkbox"/> Black cottonwood</li> <li><input type="checkbox"/> Red alder</li> <li><input type="checkbox"/> Salmonberry</li> <li><input type="checkbox"/> Nootka rose <i>Oregon ash</i></li> <li><input type="checkbox"/> Maidenhair and lady fern</li> <li><input type="checkbox"/> Blackberries</li> <li><input type="checkbox"/> Dune grasses <i>reed canary grass</i></li> </ul>	<ul style="list-style-type: none"> <li><input type="checkbox"/> Sediment lines on vegetation or other fixed objects</li> <li><input checked="" type="checkbox"/> Change from channel deposits to older alluvium.</li> <li><input type="checkbox"/> Darker stain lines on fixed objects</li> <li><input type="checkbox"/> Exposed roots/root scour.</li> <li><input type="checkbox"/> Drainage patterns, as evidenced by flattened vegetation</li> <li><input type="checkbox"/> Weathered and buried driftwood</li> </ul>
Above OHWM	<ul style="list-style-type: none"> <li><input checked="" type="checkbox"/> Hillslope toe</li> <li><input checked="" type="checkbox"/> Terraces or alluvium with an organic horizon or other developed soil horizons</li> <li><input type="checkbox"/> Relic floodplain surface</li> <li><input checked="" type="checkbox"/> Well developed soil A and B horizons/duff layer</li> </ul>	<ul style="list-style-type: none"> <li><input type="checkbox"/> Indian plum</li> <li><input type="checkbox"/> Red alder</li> <li><input type="checkbox"/> Western red cedar</li> <li><input type="checkbox"/> Douglas fir</li> <li><input type="checkbox"/> Western hemlock</li> <li><input type="checkbox"/> Ponderosa pine</li> <li><input checked="" type="checkbox"/> Oregon white oak</li> <li><input type="checkbox"/> Coast pine <i>Oregon ash</i></li> <li><input type="checkbox"/> Quaking aspen <i>ash</i></li> <li><input type="checkbox"/> Vine maple (lakes)</li> <li><input type="checkbox"/> Blackberries</li> </ul>	<ul style="list-style-type: none"> <li><input type="checkbox"/> Lighter or no staining on fixed objects</li> <li><input type="checkbox"/> Overbank deposits</li> </ul>

### Notes

There is vegetation below the OHWM such as reed canary grass and needle spikegrass. The edge of the OHWM has a clear area of less vegetation




Plant Distribution Across OHWM Gradient						
Below <del>Above</del> OHWM			At/Straddling OHWM		Above OHWM	
reed canarygrass	FACW		Oregon ash	FACW	Oregon ash	FACW
needle spikenush	ORL		reed canarygrass	PPCW	Oregon white oak	FACW

20 Nov 2019  
Photo Direction: E  
WSEL: 7.68ft NAVD88

GC-D-(1-3)



Change in vegetation type from reed canary grass to upland vegetation



# Appendix A: Field data form

## General Information

Site/Project: Wapato Valley  
 Name/Owner: Plas Newydd Farm  
 Location: Gree Creek  
 Description: 45.848399, -122.777329  
points: GC-E-(1-3)

The following field form is for use in the field to help in making ordinary high water mark delineations on streams. The form should be used as a guide. A team consisting of a hydrologist/ geomorphologist and a biologist may be needed to accurately determine the ordinary high water mark.

## General Observations: Day of Site Visit


Date of site visit:	12-3-2019		
Time of site visit:	10:00		
Weather conditions:	full sun		
Watershed development:	Highly developed <input checked="" type="checkbox"/>	Mod. Developed <input type="checkbox"/>	Undeveloped <input type="checkbox"/>
Reach development:	Highly developed <input type="checkbox"/>	Mod. Developed <input checked="" type="checkbox"/>	Undeveloped <input type="checkbox"/>
Recent site disturbance?	No <input checked="" type="checkbox"/>	Yes <input type="checkbox"/>	Describe:
Upstream flow control devices?	No <input type="checkbox"/>	Yes <input checked="" type="checkbox"/>	Describe:
Bank armoring at the site?	No <input checked="" type="checkbox"/>	Yes <input type="checkbox"/>	Describe:
Bank armoring up or downstream?	No <input type="checkbox"/>	Yes <input checked="" type="checkbox"/>	Describe:
Observable tidal backwater?	No <input type="checkbox"/>	Yes <input checked="" type="checkbox"/>	
In-water structures? (i.e. bridge pilings, railroad embankments)	No <input type="checkbox"/>	Yes <input checked="" type="checkbox"/>	Describe: <u>pilings in Columbia River</u>
Animals grazing in riparian zone?	No <input checked="" type="checkbox"/>	Yes <input type="checkbox"/>	Describe:
Observable beaver activity?	No <input checked="" type="checkbox"/>	Yes <input type="checkbox"/>	Describe:

## Complete Vegetation Transects

- Use guidelines in Chapter 4 to complete vegetation transects.
- Determine upper and lower bounds of the OHWM from vegetation transects.
- After completing vegetation transects, look for more field indicators near the upper and lower bounds of the OHWM. Use the checklist as guidance.

## Sketch

If a simple site, sketch a cross-sectional diagram of the site below. Include location of the waterway and upper and lower bounds of the OHWM defined by the vegetation communities or other OHWM indicators. Page 3 of the data form can be used for more complex sketches



## Additional Indicators

Check the indicators that are observable at the site that provide rationale for establishing the OHWM at this location. The rationale should be described in detail in the report and should be supported with photographs taken during the site visit.

	Soil and geomorphic indicators <sup>24</sup>	Vegetative indicators <sup>25</sup>	Other indicators
<b>Below OHWM</b>	<ul style="list-style-type: none"> <li>○ Sediment bars</li> <li>○ Scour line</li> <li>○ Clean cobbles/boulders.</li> <li>☒ Bank erosion/scour</li> <li>○ Lack of soil horizons</li> </ul>	Vegetation tolerant of inundation or high flow disturbances such as: <ul style="list-style-type: none"> <li>○ Willows</li> <li>○ Black cottonwood</li> <li>○ Japanese knotweed</li> <li>○ Skunk cabbage</li> <li>○ Aquatic plants</li> </ul>	<ul style="list-style-type: none"> <li>○ Exposed roots/root scour</li> <li>☒ Drainage patterns, as shown by flattened vegetation.</li> <li>☒ Aquatic animals</li> <li>○ Algal mats</li> <li>○ Iron staining</li> </ul>

<sup>24</sup> Refer to Chapter 4 for a more complete description of indicators.

<sup>25</sup> Species are provided as examples. Refer to Appendix B for a more complete listing of plant species and their distribution across the OHWM gradient. Some species occur in more than one category depending on site conditions. For example Indian plum and red alder may straddle the OHWM where soil drainage is high. They may occur above OHWM where soil drainage is low to moderate.

	Soil and geomorphic indicators <sup>24</sup>	Vegetative indicators <sup>25</sup>	Other indicators
At or straddling OHWM	<ul style="list-style-type: none"> <li>○ Top of bank</li> <li>○ Toe of lowest terrace (if terrace has developed horizons which may include a duff layer and A and B horizons versus freshly deposited alluvium)</li> <li>☒ Benches</li> </ul>	<ul style="list-style-type: none"> <li>○ Willows</li> <li>○ Western red cedar</li> <li>○ Vine maple (streams)</li> <li>○ Black cottonwood</li> <li>○ Red alder <i>CAVEY</i></li> <li>○ Salmonberry <i>obscure</i></li> <li>○ Nootka rose</li> <li>○ Maidenhair and lady fern</li> <li>○ Blackberries</li> <li>○ Dunegrasses</li> </ul>	<ul style="list-style-type: none"> <li>○ Sediment lines on vegetation or other fixed objects</li> <li>○ Change from channel deposits to older alluvium.</li> <li>○ Darker stain lines on fixed objects</li> <li>○ Exposed roots/root scour.</li> <li>○ Drainage patterns, as evidenced by flattened vegetation</li> <li>☒ Weathered and buried driftwood</li> </ul>
Above OHWM	<ul style="list-style-type: none"> <li>○ Hillslope toe</li> <li>☒ Terraces or alluvium with an organic horizon or other developed soil horizons</li> <li>☒ Relic floodplain surface</li> <li>☒ Well developed soil A and B horizons/duff layer</li> </ul>	<ul style="list-style-type: none"> <li>○ Indian plum <i>OREGON</i></li> <li>○ Red alder <i>ASH</i></li> <li>○ Western red cedar</li> <li>○ Douglas fir</li> <li>○ Western hemlock</li> <li>○ Ponderosa pine</li> <li>○ Oregon white oak</li> <li>○ Coast pine</li> <li>○ Quaking aspen</li> <li>○ Vine maple (lakes)</li> <li>☒ Blackberries</li> </ul>	<ul style="list-style-type: none"> <li>☒ Lighter or no staining on fixed objects</li> <li>○ Overbank deposits</li> </ul>

Notes

---



---



---



---



---



---



---




---



---



---



Below <del>Above</del> OHWM	Plant Distribution Across OHWM Gradient			Above OHWM
	At/Straddling OHWM			
none	slough sedge	OBL	Oregon ash	FACW
	reed canarygrass	FACW	himalaya blackberry	FACW
			reed canarygrass	FACW
			red-osier dogwood	FACW

3 Dec 2019  
Photo Direction: S  
WSEL: 8.39ft NAVD88

GC-E-(1-3)



← OHWM: Wrack line



# Appendix A: Field data form

## General Information

Site/Project  
 Name/Owner: Wapato Valley/Plas Newydd LLC  
 Location: Lancaster Lake  
 Description: 45.848966, -122.751122  
points: LL-A-(1-4)

The following field form is for use in the field to help in making ordinary high water mark delineations on streams. The form should be used as a guide. A team consisting of a hydrologist/ geomorphologist and a biologist may be needed to accurately determine the ordinary high water mark.

## General Observations: Day of Site Visit


Date of site visit:	Nov 19 2019		
Time of site visit:	11:23		
Weather conditions:	Over cast		
Watershed development:	Highly developed <input checked="" type="checkbox"/>	Mod. Developed <input type="checkbox"/>	Undeveloped <input type="checkbox"/>
Reach development:	Highly developed <input type="checkbox"/>	Mod. Developed <input checked="" type="checkbox"/>	Undeveloped <input type="checkbox"/>
Recent site disturbance?	No <input checked="" type="checkbox"/>	Yes <input type="checkbox"/>	Describe:
Upstream flow control devices?	No <input type="checkbox"/>	Yes <input checked="" type="checkbox"/>	Describe: levee with tide gate
Bank armoring at the site?	No <input type="checkbox"/>	Yes <input checked="" type="checkbox"/>	Describe: levee
Bank armoring up or downstream?	No <input checked="" type="checkbox"/>	Yes <input type="checkbox"/>	Describe: lake is impounded
Observable tidal backwater?	No <input checked="" type="checkbox"/>	Yes <input type="checkbox"/>	
In-water structures? (i.e. bridge pilings, railroad embankments)	No <input type="checkbox"/>	Yes <input checked="" type="checkbox"/>	Describe: levee and tide gate
Animals grazing in riparian zone?	No <input checked="" type="checkbox"/>	Yes <input type="checkbox"/>	Describe:
Observable beaver activity?	No <input type="checkbox"/>	Yes <input checked="" type="checkbox"/>	Describe: Beaver channels

## Complete Vegetation Transects

- Use guidelines in Chapter 4 to complete vegetation transects.
- Determine upper and lower bounds of the OHWM from vegetation transects.
- After completing vegetation transects, look for more field indicators near the upper and lower bounds of the OHWM. Use the checklist as guidance.

## Sketch

If a simple site, sketch a cross-sectional diagram of the site below. Include location of the waterway and upper and lower bounds of the OHWM defined by the vegetation communities or other OHWM indicators. Page 3 of the data form can be used for more complex sketches



## Additional Indicators

Check the indicators that are observable at the site that provide rationale for establishing the OHWM at this location. The rationale should be described in detail in the report and should be supported with photographs taken during the site visit.

	Soil and geomorphic indicators <sup>24</sup>	Vegetative indicators <sup>25</sup>	Other indicators
<b>Below OHWM</b>	<ul style="list-style-type: none"> <li>○ Sediment bars</li> <li>○ Scour line</li> <li>○ Clean cobbles/boulders.</li> <li>○ Bank erosion/scour</li> <li>○ Lack of soil horizons</li> </ul>	Vegetation tolerant of inundation or high flow disturbances such as: <ul style="list-style-type: none"> <li>○ Willows</li> <li>○ Black cottonwood</li> <li>○ Japanese knotweed</li> <li>○ Skunk cabbage</li> <li>○ Aquatic plants</li> </ul>	<ul style="list-style-type: none"> <li>○ Exposed roots/root scour</li> <li>○ Drainage patterns, as shown by flattened vegetation.</li> <li><input checked="" type="checkbox"/> Aquatic animals</li> <li>○ Algal mats</li> <li>○ Iron staining</li> </ul>


<sup>24</sup> Refer to Chapter 4 for a more complete description of indicators.

<sup>25</sup> Species are provided as examples. Refer to Appendix B for a more complete listing of plant species and their distribution across the OHWM gradient. Some species occur in more than one category depending on site conditions. For example Indian plum and red alder may straddle the OHWM where soil drainage is high. They may occur above OHWM where soil drainage is low to moderate.

	Soil and geomorphic indicators <sup>24</sup>	Vegetative indicators <sup>25</sup>	Other indicators
At or straddling OHWM	<ul style="list-style-type: none"> <li>○ Top of bank</li> <li>☒ Toe of lowest terrace (if terrace has developed horizons which may include a duff layer and A and B horizons versus freshly deposited alluvium)</li> <li>○ Benches</li> </ul>	<ul style="list-style-type: none"> <li>○ Willows</li> <li>○ Western red cedar</li> <li>○ Vine maple (streams)</li> <li>○ Black cottonwood</li> <li>○ Red alder</li> <li>○ Salmonberry</li> <li>○ Nootka rose</li> <li>○ Maidenhair and lady fern</li> <li>○ Blackberries</li> <li>○ Dune grasses <i>reed canary grass</i></li> </ul>	<ul style="list-style-type: none"> <li>☒ Sediment lines on vegetation or other fixed objects</li> <li>○ Change from channel deposits to older alluvium.</li> <li>☒ Darker stain lines on fixed objects</li> <li>○ Exposed roots/root scour.</li> <li>○ Drainage patterns, as evidenced by flattened vegetation</li> <li>○ Weathered and buried driftwood</li> </ul>
Above OHWM	<ul style="list-style-type: none"> <li>☒ Hillslope toe</li> <li>☒ Terraces or alluvium with an organic horizon or other developed soil horizons</li> <li>○ Relic floodplain surface</li> <li>☒ Well developed soil A and B horizons/duff layer</li> </ul>	<ul style="list-style-type: none"> <li>○ Indian plum</li> <li>○ Red alder</li> <li>○ Western red cedar</li> <li>☒ Douglas fir</li> <li>○ Western hemlock</li> <li>○ Ponderosa pine</li> <li>☒ Oregon white oak</li> <li>○ Coast pine</li> <li>○ Quaking aspen</li> <li>☒ Vine maple (lakes)</li> <li>☒ Blackberries</li> </ul>	<ul style="list-style-type: none"> <li>○ Lighter or no staining on fixed objects</li> <li>○ Overbank deposits</li> </ul>

Notes

Impounded lake. A tree fell over and there is sediment staining at OHWM. Vegetation consists of reed canary grass and bull rush



Plant Distribution Across OHWM Gradient					
Below <del>Above</del> OHWM		At/Straddling OHWM		Above OHWM	
reed canarygrass	FACW	reed canarygrass	FACW	Oregon white oak	FACU
slough sedge	OBL			Oregon ash	FACW
				Scott's broom	NI
				licorice fern	NI

19 Nov 2019

Photo Direction: S

WSEL: 8.70ft NAVD88

LL-A-(1-4)



← OHWM: Sediment line on tree

# Appendix A: Field data form

## General Information

Site/Project  
 Name/Owner: Wapato Valley/Plas Nwidd LLC  
 Location: Lancaster Lake  
 Description: 45.850023 -122.750232  
points: LL-B-(1-8)

The following field form is for use in the field to help in making ordinary high water mark delineations on streams. The form should be used as a guide. A team consisting of a hydrologist/ geomorphologist and a biologist may be needed to accurately determine the ordinary high water mark.

## General Observations: Day of Site Visit


Date of site visit:	Nov 19 2019		
Time of site visit:	10:43		
Weather conditions:	overcast		
Watershed development:	Highly developed <input checked="" type="radio"/>	Mod. Developed <input type="radio"/>	Undeveloped <input type="radio"/>
Reach development:	Highly developed <input type="radio"/>	Mod. Developed <input checked="" type="radio"/>	Undeveloped <input type="radio"/>
Recent site disturbance?	No <input checked="" type="radio"/>	Yes <input type="radio"/>	Describe:
Upstream flow control devices?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: tide gate
Bank armoring at the site?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: levee with tide gate
Bank armoring up or downstream?	No <input checked="" type="radio"/>	Yes <input type="radio"/>	Describe: lake is impounded
Observable tidal backwater?	No <input checked="" type="radio"/>	Yes <input type="radio"/>	
In-water structures? (i.e. bridge pilings, railroad embankments)	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: levee and tide gate
Animals grazing in riparian zone?	No <input checked="" type="radio"/>	Yes <input type="radio"/>	Describe:
Observable beaver activity?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: Beaver channel

## Complete Vegetation Transects

- Use guidelines in Chapter 4 to complete vegetation transects.
- Determine upper and lower bounds of the OHWM from vegetation transects.
- After completing vegetation transects, look for more field indicators near the upper and lower bounds of the OHWM. Use the checklist as guidance.

## Sketch

If a simple site, sketch a cross-sectional diagram of the site below. Include location of the waterway and upper and lower bounds of the OHWM defined by the vegetation communities or other OHWM indicators. Page 3 of the data form can be used for more complex sketches



willows are site OHWM indicators

reed canary grass

## Additional Indicators

Check the indicators that are observable at the site that provide rationale for establishing the OHWM at this location. The rationale should be described in detail in the report and should be supported with photographs taken during the site visit.

	Soil and geomorphic indicators <sup>24</sup>	Vegetative indicators <sup>25</sup>	Other indicators
<b>Below OHWM</b>	<ul style="list-style-type: none"> <li>○ Sediment bars</li> <li>○ Scour line</li> <li>○ Clean cobbles/boulders.</li> <li>○ Bank erosion/scour</li> <li>○ Lack of soil horizons</li> </ul>	Vegetation tolerant of inundation or high flow disturbances such as: <ul style="list-style-type: none"> <li>○ Willows</li> <li>○ Black cottonwood</li> <li>○ Japanese knotweed</li> <li>○ Skunk cabbage</li> <li>⊗ Aquatic plants</li> </ul>	<ul style="list-style-type: none"> <li>○ Exposed roots/root scour</li> <li>○ Drainage patterns, as shown by flattened vegetation.</li> <li>⊗ Aquatic animals</li> <li>○ Algal mats</li> <li>○ Iron staining</li> </ul>

<sup>24</sup> Refer to Chapter 4 for a more complete description of indicators.


<sup>25</sup> Species are provided as examples. Refer to Appendix B for a more complete listing of plant species and their distribution across the OHWM gradient. Some species occur in more than one category depending on site conditions. For example Indian plum and red alder may straddle the OHWM where soil drainage is high. They may occur above OHWM where soil drainage is low to moderate.

	Soil and geomorphic indicators <sup>24</sup>	Vegetative indicators <sup>25</sup>	Other indicators
At or straddling OHWM	<ul style="list-style-type: none"> <li>○ Top of bank</li> <li>○ Toe of lowest terrace (if terrace has developed horizons which may include a duff layer and A and B horizons versus freshly deposited alluvium)</li> <li>○ Benches</li> </ul>	<ul style="list-style-type: none"> <li>⊗ Willows</li> <li>○ Western red cedar</li> <li>○ Vine maple (streams)</li> <li>○ Black cottonwood</li> <li>○ Red alder</li> <li>○ Salmonberry</li> <li>○ Nootka rose</li> <li>○ Maidenhair and lady fern</li> <li>○ Blackberries</li> <li>○ Dunegrasses</li> </ul>	<ul style="list-style-type: none"> <li>○ Sediment lines on vegetation or other fixed objects</li> <li>○ Change from channel deposits to older alluvium.</li> <li>○ Darker stain lines on fixed objects</li> <li>○ Exposed roots/root scour.</li> <li>○ Drainage patterns, as evidenced by flattened vegetation</li> <li>○ Weathered and buried driftwood</li> </ul>
Above OHWM	<ul style="list-style-type: none"> <li>⊗ Hillslope toe</li> <li>○ Terraces or alluvium with an organic horizon or other developed soil horizons</li> <li>⊗ Relic floodplain surface</li> <li>⊗ Well developed soil A and B horizons/duff layer</li> </ul>	<ul style="list-style-type: none"> <li>○ Indian plum</li> <li>○ Red alder</li> <li>○ Western red cedar</li> <li>⊗ Douglas fir</li> <li>○ Western hemlock</li> <li>○ Ponderosa pine</li> <li>⊗ Oregon white oak</li> <li>○ Coast pine</li> <li>○ Quaking aspen</li> <li>⊗ Vine maple (lakes)</li> <li>⊗ Blackberries</li> </ul>	<ul style="list-style-type: none"> <li>○ Lighter or no staining on fixed objects</li> <li>○ Overbank deposits</li> </ul>

### Notes

This is an impounded lake. There is a break in vegetation habitats. Polygonums is at lower elevation and reed canary grass and willows are at higher elevations indicating OHWM





Plant Distribution Across OHWM Gradient					
Below <del>Above</del> OHWM		At/Straddling OHWM		Above OHWM	
spotted lady's thumb	FACW	reed canarygrass	FACW	willows (mature)	FACW
nodding beggarstick	OBL	willow (sprouts)	FACW	reed canarygrass	FACW
wapato	OBL				

19 Nov 2019

Photo Direction: N

WSEL: 8.84ft NAVD88

LL-B-(1-8)

OHWL: Change in vegetation type  
from polygonum species to willow



# Appendix A: Field data form

## General Information

Site/Project: Wapato Valley /  
 Name/Owner: Plas Newydd Farm  
 Location: Lancaster Lake  
 Description: 45.85181, -122.748098  
points: LL-C-(1-6)

The following field form is for use in the field to help in making ordinary high water mark delineations on streams. The form should be used as a guide. A team consisting of a hydrologist/ geomorphologist and a biologist may be needed to accurately determine the ordinary high water mark.

## General Observations: Day of Site Visit


Date of site visit:	11-19-2019		
Time of site visit:	11:05		
Weather conditions:	overcast		
Watershed development:	Highly developed <input checked="" type="radio"/>	Mod. Developed <input type="radio"/>	Undeveloped <input type="radio"/>
Reach development:	Highly developed <input type="radio"/>	Mod. Developed <input checked="" type="radio"/>	Undeveloped <input type="radio"/>
Recent site disturbance?	No <input checked="" type="radio"/>	Yes <input type="radio"/>	Describe:
Upstream flow control devices?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: Levee w/ tidegate water control structure that block tidal influence.
Bank armoring at the site?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: Levee is armored.
Bank armoring up or downstream?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: Levee and parts of Cree Creek upstream
Observable tidal backwater?	No <input checked="" type="radio"/>	Yes <input type="radio"/>	
In-water structures? (i.e. bridge pilings, railroad embankments)	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: Levee w/ tidegate
Animals grazing in riparian zone?	No <input checked="" type="radio"/>	Yes <input type="radio"/>	Describe:
Observable beaver activity?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: Several channels, lodges and a small dam.

## Complete Vegetation Transects

- Use guidelines in Chapter 4 to complete vegetation transects.
- Determine upper and lower bounds of the OHWM from vegetation transects.
- After completing vegetation transects, look for more field indicators near the upper and lower bounds of the OHWM. Use the checklist as guidance.

## Sketch

If a simple site, sketch a cross-sectional diagram of the site below. Include location of the waterway and upper and lower bounds of the OHWM defined by the vegetation communities or other OHWM indicators. Page 3 of the data form can be used for more complex sketches



## Additional Indicators

Check the indicators that are observable at the site that provide rationale for establishing the OHWM at this location. The rationale should be described in detail in the report and should be supported with photographs taken during the site visit.

	Soil and geomorphic indicators <sup>24</sup>	Vegetative indicators <sup>25</sup>	Other indicators
<b>Below OHWM</b>	<ul style="list-style-type: none"> <li>○ Sediment bars</li> <li>○ Scour line</li> <li>○ Clean cobbles/boulders.</li> <li>○ Bank erosion/scour</li> <li>○ Lack of soil horizons</li> </ul>	Vegetation tolerant of inundation or high flow disturbances such as: <ul style="list-style-type: none"> <li>○ Willows</li> <li>○ Black cottonwood</li> <li>○ Japanese knotweed + other</li> <li>○ Skunk cabbage</li> <li>○ Aquatic plants</li> </ul>	<ul style="list-style-type: none"> <li>○ Exposed roots/root scour</li> <li>○ Drainage patterns, as shown by flattened vegetation.</li> <li>○ Aquatic animals</li> <li>○ Algal mats</li> <li>○ Iron staining</li> </ul>


<sup>24</sup> Refer to Chapter 4 for a more complete description of indicators.

<sup>25</sup> Species are provided as examples. Refer to Appendix B for a more complete listing of plant species and their distribution across the OHWM gradient. Some species occur in more than one category depending on site conditions. For example Indian plum and red alder may straddle the OHWM where soil drainage is high. They may occur above OHWM where soil drainage is low to moderate.

	Soil and geomorphic indicators <sup>24</sup>	Vegetative indicators <sup>25</sup>	Other indicators
At or straddling OHWM	<ul style="list-style-type: none"> <li>○ Top of bank</li> <li>○ Toe of lowest terrace (if terrace has developed horizons which may include a duff layer and A and B horizons versus freshly deposited alluvium)</li> <li>○ Benches</li> </ul>	<ul style="list-style-type: none"> <li>☒ Willows</li> <li>○ Western red cedar</li> <li>○ Vine maple (streams)</li> <li>○ Black cottonwood</li> <li>○ Red alder <i>Red</i></li> <li>○ Salmonberry <i>Cowberry</i></li> <li>○ Nootka rose</li> <li>○ Maidenhair and lady fern</li> <li>○ Blackberries</li> <li>○ Dunegrasses</li> </ul>	<ul style="list-style-type: none"> <li>○ Sediment lines on vegetation or other fixed objects</li> <li>○ Change from channel deposits to older alluvium.</li> <li>○ Darker stain lines on fixed objects</li> <li>○ Exposed roots/root scour.</li> <li>○ Drainage patterns, as evidenced by flattened vegetation</li> <li>○ Weathered and buried driftwood</li> </ul>
Above OHWM	<ul style="list-style-type: none"> <li>☒ Hillslope toe</li> <li>○ Terraces or alluvium with an organic horizon or other developed soil horizons</li> <li>○ Relic floodplain surface</li> <li>☒ Well developed soil A and B horizons/duff layer</li> </ul>	<ul style="list-style-type: none"> <li>○ Indian plum</li> <li>○ Red alder <i>Oregon Ash</i></li> <li>○ Western red cedar</li> <li>○ Douglas fir</li> <li>○ Western hemlock</li> <li>○ Ponderosa pine</li> <li>☒ Oregon white oak</li> <li>○ Coast pine</li> <li>○ Quaking aspen</li> <li>☒ Vine maple (lakes)</li> <li>○ Blackberries</li> </ul>	<ul style="list-style-type: none"> <li>○ Lighter or no staining on fixed objects</li> <li>○ Overbank deposits</li> </ul>

## Notes

Lancaster Lake is impounded at its southern end where it drains into Bee Creek but is blocked from tidal influence. Beaver have created many channels, lodges and a dam. Water from adjacent farm fields drains into the lake via ditches on the northeast and northwest points of the lake.



Plant Distribution Across OHWM Gradient					
Below <del>Above</del> OHWM		At/Straddling OHWM		Above OHWM	
spotted ladythumb	FACW	reed canarygrass	FACW	willow (mature)	FACW
nodding beargrass	OBL	willow (sprouts)	FACW	reed canarygrass	FACW
wapato	OBL				

19 Nov 2019  
Photo Direction: E  
WSEL: 8.72ft NAVD88

LL-C-(1-6)



← OHWM: Change in vegetation type from polygonum species to willow

LL

## Appendix A: Field data form

### General Information

Site/Project: \_\_\_\_\_  
 Name/Owner: Wapato Valley / Plas Newydd Farm  
 Location: Lancaster Lake  
 Description: 45.849093, -122.750488  
 points: LL-D-(1-4)

The following field form is for use in the field to help in making ordinary high water mark delineations on streams. The form should be used as a guide. A team consisting of a hydrologist/ geomorphologist and a biologist may be needed to accurately determine the ordinary high water mark.

### General Observations: Day of Site Visit

Date of site visit:	12-4-2019		
Time of site visit:	11:30		
Weather conditions:	sun		
Watershed development:	Highly developed <input type="radio"/>	Mod. Developed <input type="radio"/>	Undeveloped <input type="radio"/>
Reach development:	Highly developed <input type="radio"/>	Mod. Developed <input type="radio"/>	Undeveloped <input type="radio"/>
Recent site disturbance?	No <input checked="" type="radio"/>	Yes <input type="radio"/>	Describe:
Upstream flow control devices?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: tidegate between lake and Cree Creek
Bank armoring at the site?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: levee w/ tidegate
Bank armoring up or downstream?	No <input checked="" type="radio"/>	Yes <input type="radio"/>	Describe: lake is impounded
Observable tidal backwater?	No <input checked="" type="radio"/>	Yes <input type="radio"/>	
In-water structures? (i.e. bridge pilings, railroad embankments)	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: levee and tidegate
Animals grazing in riparian zone?	No <input checked="" type="radio"/>	Yes <input type="radio"/>	Describe:
Observable beaver activity?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: beaver lodges and channels

### Complete Vegetation Transects


- Use guidelines in Chapter 4 to complete vegetation transects.
- Determine upper and lower bounds of the OHWM from vegetation transects.
- After completing vegetation transects, look for more field indicators near the upper and lower bounds of the OHWM. Use the checklist as guidance.



LL

### Sketch

If a simple site, sketch a cross-sectional diagram of the site below. Include location of the waterway and upper and lower bounds of the OHWM defined by the vegetation communities or other OHWM indicators. Page 3 of the data form can be used for more complex sketches



### Additional Indicators

Check the indicators that are observable at the site that provide rationale for establishing the OHWM at this location. The rationale should be described in detail in the report and should be supported with photographs taken during the site visit.

	Soil and geomorphic indicators <sup>24</sup>	Vegetative indicators <sup>25</sup>	Other indicators
<b>Below OHWM</b>	<ul style="list-style-type: none"> <li><input type="radio"/> Sediment bars</li> <li><input type="radio"/> Scour line</li> <li><input checked="" type="checkbox"/> Clean cobbles/boulders.</li> <li><input type="radio"/> Bank erosion/scour</li> <li><input checked="" type="checkbox"/> Lack of soil horizons</li> </ul>	Vegetation tolerant of inundation or high flow disturbances such as: <ul style="list-style-type: none"> <li><input type="radio"/> Willows</li> <li><input type="radio"/> Black cottonwood</li> <li><input type="radio"/> Japanese knotweed</li> <li><input type="radio"/> Skunk cabbage</li> <li><input checked="" type="checkbox"/> Aquatic plants</li> </ul>	<ul style="list-style-type: none"> <li><input type="radio"/> Exposed roots/root scour</li> <li><input type="radio"/> Drainage patterns, as shown by flattened vegetation</li> <li><input checked="" type="checkbox"/> Aquatic animals</li> <li><input type="radio"/> Algal mats</li> <li><input type="radio"/> Iron staining</li> </ul>

<sup>24</sup> Refer to Chapter 4 for a more complete description of indicators.

<sup>25</sup> Species are provided as examples. Refer to Appendix B for a more complete listing of plant species and their distribution across the OHWM gradient. Some species occur in more than one category depending on site conditions. For example Indian plum and red alder may straddle the OHWM where soil drainage is high. They may occur above OHWM where soil drainage is low to moderate.

LL

	Soil and geomorphic indicators <sup>24</sup>	Vegetative indicators <sup>25</sup>	Other indicators
At or straddling OHWM	<ul style="list-style-type: none"> <li>o Top of bank</li> <li>o Toe of lowest terrace (if terrace has developed horizons which may include a duff layer and A and B horizons versus freshly deposited alluvium)</li> <li>o Benches</li> </ul>	<ul style="list-style-type: none"> <li>o Willows</li> <li>o Western red cedar</li> <li>o Vine maple (streams)</li> <li>o Black cottonwood</li> <li>o Red alder <i>o reed</i></li> <li>o Salmonberry <i>canarygrass</i></li> <li>o Nootka rose</li> <li>o Maidenhair and lady fern</li> <li>o Blackberries</li> <li>o Dunegrasses</li> </ul>	<ul style="list-style-type: none"> <li>o Sediment lines on vegetation or other fixed objects</li> <li>o Change from channel deposits to older alluvium.</li> <li>o Darker stain lines on fixed objects</li> <li>o Exposed roots/root scour.</li> <li>o Drainage patterns, as evidenced by flattened vegetation</li> <li>o Weathered and buried driftwood</li> </ul>
Above OHWM	<ul style="list-style-type: none"> <li>o Hillslope toe</li> <li>o Terraces or alluvium with an organic horizon or other developed soil horizons</li> <li>o Relic floodplain surface</li> <li>o Well developed soil A and B horizons/duff layer</li> </ul>	<ul style="list-style-type: none"> <li>o Indian plum <i>o Scott's</i></li> <li>o Red alder <i>broom</i></li> <li>o Western red cedar</li> <li>o Douglas fir</li> <li>o Western hemlock <i>o cany</i></li> <li>o Ponderosa pine</li> <li>o Oregon white oak</li> <li>o Coast pine</li> <li>o Quaking aspen</li> <li>o Vine maple (lakes)</li> <li>o Blackberries</li> </ul>	<ul style="list-style-type: none"> <li>o Lighter or no staining on fixed objects</li> <li>o Overbank deposits</li> </ul>

Notes

The best indicator here is the break between more water tolerant reed canarygrass and upland Scott's broom.

---



---



---




---



---

LL



Plant Distribution Across OHWM Gradient					
Below <del>Above</del> OHWM		At/Straddling OHWM		Above OHWM	
rough cocklebur	FAC	reed canarygrass	FACW	Scott's broom	NI
reed canarygrass	FACW	Robert geranium	FACU	queen Anne's lace	UPL
		birdsfoot trefoil	FACU	licorice fern	NI
		Scott's broom	NI	Oregon white oak	FACU
				Douglas fir	FACU
				oxeye daisy	FACU
				camas	FACW
				snowberry	FACU
				shiny geranium	NI
				barnyard grass	FAC


4 Dec 2019

Photo Direction: S

WSEL: 8.84ft NAVD88

LL-D-(1-4)

OHWM: Change in vegetation from water tolerant species to upland vegetation



# Appendix A: Field data form

## General Information

Site/Project: Wapato Valley  
 Name/Owner: Plas Newyald Farm  
 Location: Lewis River  
 Description: 45.863632, -122.750108  
points: LR-A-(1-2)

The following field form is for use in the field to help in making ordinary high water mark delineations on streams. The form should be used as a guide. A team consisting of a hydrologist/ geomorphologist and a biologist may be needed to accurately determine the ordinary high water mark.

## General Observations: Day of Site Visit


Date of site visit:	<u>20 Nov 2014</u>		
Time of site visit:	<u>15:39</u>		
Weather conditions:	<u>Full Sun</u>		
Watershed development:	Highly developed <input checked="" type="radio"/>	Mod. Developed <input type="radio"/>	Undeveloped <input type="radio"/>
Reach development:	Highly developed <input type="radio"/>	Mod. Developed <input type="radio"/>	Undeveloped <input type="radio"/>
Recent site disturbance?	No <input checked="" type="radio"/>	Yes <input type="radio"/>	Describe:
Upstream flow control devices?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: <u>Mervin Dam/Bonneville Dam</u>
Bank armoring at the site?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: <u>rock armoring</u>
Bank armoring up or downstream?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: <u>Both up and down stream</u>
Observable tidal backwater?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	
In-water structures? (i.e. bridge pilings, railroad embankments)	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: <u>Railroad embankments/pilings</u>
Animals grazing in riparian zone?	No <input checked="" type="radio"/>	Yes <input type="radio"/>	Describe:
Observable beaver activity?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: <u>beaver chews</u>

## Complete Vegetation Transects

- Use guidelines in Chapter 4 to complete vegetation transects.
- Determine upper and lower bounds of the OHWM from vegetation transects.
- After completing vegetation transects, look for more field indicators near the upper and lower bounds of the OHWM. Use the checklist as guidance.

## Sketch

If a simple site, sketch a cross-sectional diagram of the site below. Include location of the waterway and upper and lower bounds of the OHWM defined by the vegetation communities or other OHWM indicators. Page 3 of the data form can be used for more complex sketches



## Additional Indicators

Check the indicators that are observable at the site that provide rationale for establishing the OHWM at this location. The rationale should be described in detail in the report and should be supported with photographs taken during the site visit.

	Soil and geomorphic indicators <sup>24</sup>	Vegetative indicators <sup>25</sup>	Other indicators
Below OHWM	<ul style="list-style-type: none"> <li>○ Sediment bars</li> <li>○ Scour line</li> <li>○ Clean cobbles/boulders.</li> <li>○ Bank erosion/scour</li> <li><input checked="" type="checkbox"/> Lack of soil horizons</li> </ul>	Vegetation tolerant of inundation or high flow disturbances such as: <ul style="list-style-type: none"> <li>○ Willows</li> <li>○ Black cottonwood</li> <li>○ Japanese knotweed</li> <li>○ Skunk cabbage</li> <li><input checked="" type="checkbox"/> Aquatic plants</li> </ul>	<ul style="list-style-type: none"> <li>○ Exposed roots/root scour</li> <li>○ Drainage patterns, as shown by flattened vegetation.</li> <li><input checked="" type="checkbox"/> Aquatic animals</li> <li>○ Algal mats</li> <li>○ Iron staining</li> </ul>


<sup>24</sup> Refer to Chapter 4 for a more complete description of indicators.

<sup>25</sup> Species are provided as examples. Refer to Appendix B for a more complete listing of plant species and their distribution across the OHWM gradient. Some species occur in more than one category depending on site conditions. For example Indian plum and red alder may straddle the OHWM where soil drainage is high. They may occur above OHWM where soil drainage is low to moderate.

	Soil and geomorphic indicators <sup>24</sup>	Vegetative indicators <sup>25</sup>	Other indicators
At or straddling OHWM	<ul style="list-style-type: none"> <li>○ Top of bank</li> <li>☒ Toe of lowest terrace (if terrace has developed horizons which may include a duff layer and A and B horizons versus freshly deposited alluvium)</li> <li>○ Benches</li> </ul>	<ul style="list-style-type: none"> <li>○ Willows</li> <li>○ Western red cedar</li> <li>○ Vine maple (streams)</li> <li>○ Black cottonwood</li> <li>○ Red alder</li> <li>○ Salmonberry</li> <li>○ Nootka rose</li> <li>○ Maidenhair and lady fern</li> <li>○ Blackberries <i>red</i></li> <li>○ Dunegrasses <i>canarygrass</i></li> </ul>	<ul style="list-style-type: none"> <li>☒ Sediment lines on vegetation or other fixed objects</li> <li>☒ Change from channel deposits to older alluvium.</li> <li>☒ Darker stain lines on fixed objects</li> <li>○ Exposed roots/root scour.</li> <li>○ Drainage patterns, as evidenced by flattened vegetation</li> <li>○ Weathered and buried driftwood</li> </ul>
Above OHWM	<ul style="list-style-type: none"> <li>○ Hillislope toe</li> <li>○ Terraces or alluvium with an organic horizon or other developed soil horizons</li> <li>○ Relic floodplain surface</li> <li>○ Well developed soil A and B horizons/duff layer</li> </ul>	<ul style="list-style-type: none"> <li>○ Indian plum</li> <li>○ Red alder</li> <li>○ Western red cedar</li> <li>○ Douglas fir</li> <li>○ Western hemlock</li> <li>○ Ponderosa pine</li> <li>○ Oregon white oak</li> <li>○ Coast pine <i>black cottonwood</i></li> <li>○ Quaking aspen</li> <li>○ Vine maple (lakes)</li> <li>☒ Blackberries</li> </ul>	<ul style="list-style-type: none"> <li>○ Lighter or no staining on fixed objects</li> <li>○ Overbank deposits</li> </ul>

Notes

At this site the vegetation break along the shoreline and sediment deposits on riprap and a boat launch were used as the best OHWM indicators.



Plant Distribution Across OHWM Gradient					
Below <del>Above</del> OHWM		At/Straddling OHWM		Above OHWM	
reed canarygrass	FACW	red-osier dogwood	FACW	himalaya blackberry	FACW
slough sedge	OBL	willow sp.	FACW	black cottonwood	NI
purple loosestrife	OBL	spirea	FACW	moss sp.	NI
wooly sedge	OBL	pennyroyal	OBL	western dock	FACW
sneezeweed	FACW			bigleaf maple	FACW
				queen Anne's lace	FACW



20 Nov 2019  
Photo Direction: W  
WSEL: 7.16ft NAVD88

LR-A-(1-2)

← OHWM: Sediment  
line on rocks



## Appendix A: Field data form

### General Information

Site/Project: Wapato Valley  
 Name/Owner: Plas Newydd Farm  
 Location: Lewis River  
 Description: 45.861905, -122.757252  
points: LR-B-(1-6)

The following field form is for use in the field to help in making ordinary high water mark delineations on streams. The form should be used as a guide. A team consisting of a hydrologist/ geomorphologist and a biologist may be needed to accurately determine the ordinary high water mark.

### General Observations: Day of Site Visit


Date of site visit:	11-20-2019		
Time of site visit:	15:30		
Weather conditions:	full sun		
Watershed development:	Highly developed <input checked="" type="checkbox"/>	Mod. Developed <input type="checkbox"/>	Undeveloped <input type="checkbox"/>
Reach development:	Highly developed <input checked="" type="checkbox"/>	Mod. Developed <input type="checkbox"/>	Undeveloped <input type="checkbox"/>
Recent site disturbance?	No <input checked="" type="checkbox"/>	Yes <input type="checkbox"/>	Describe:
Upstream flow control devices?	No <input type="checkbox"/>	Yes <input checked="" type="checkbox"/>	Describe: Menwin Dam
Bank armoring at the site?	No <input type="checkbox"/>	Yes <input checked="" type="checkbox"/>	Describe: Riprap on both sides
Bank armoring up or downstream?	No <input type="checkbox"/>	Yes <input checked="" type="checkbox"/>	Describe: both sides up and down
Observable tidal backwater?	No <input type="checkbox"/>	Yes <input checked="" type="checkbox"/>	
In-water structures? (i.e. bridge pilings, railroad embankments)	No <input type="checkbox"/>	Yes <input checked="" type="checkbox"/>	Describe: railroad bridge, pilings
Animals grazing in riparian zone?	No <input checked="" type="checkbox"/>	Yes <input type="checkbox"/>	Describe:
Observable beaver activity?	No <input checked="" type="checkbox"/>	Yes <input type="checkbox"/>	Describe:

### Complete Vegetation Transects

- Use guidelines in Chapter 4 to complete vegetation transects.
- Determine upper and lower bounds of the OHWM from vegetation transects.
- After completing vegetation transects, look for more field indicators near the upper and lower bounds of the OHWM. Use the checklist as guidance.

## Sketch

If a simple site, sketch a cross-sectional diagram of the site below. Include location of the waterway and upper and lower bounds of the OHWM defined by the vegetation communities or other OHWM indicators. Page 3 of the data form can be used for more complex sketches



## Additional Indicators

Check the indicators that are observable at the site that provide rationale for establishing the OHWM at this location. The rationale should be described in detail in the report and should be supported with photographs taken during the site visit.

	Soil and geomorphic indicators <sup>24</sup>	Vegetative indicators <sup>25</sup>	Other indicators
<b>Below OHWM</b>	<ul style="list-style-type: none"> <li>○ Sediment bars</li> <li>○ Scour line</li> <li>○ Clean cobbles/boulders.</li> <li>○ Bank erosion/scour</li> <li>☑ Lack of soil horizons</li> </ul>	Vegetation tolerant of inundation or high flow disturbances such as: <ul style="list-style-type: none"> <li>○ Willows</li> <li>○ Black cottonwood</li> <li>○ Japanese knotweed</li> <li>○ Skunk cabbage</li> <li>○ Aquatic plants</li> </ul>	<ul style="list-style-type: none"> <li>○ Exposed roots/root scour</li> <li>○ Drainage patterns, as shown by flattened vegetation.</li> <li>☑ Aquatic animals</li> <li>○ Algal mats</li> <li>○ Iron staining</li> </ul>


<sup>24</sup> Refer to Chapter 4 for a more complete description of indicators.

<sup>25</sup> Species are provided as examples. Refer to Appendix B for a more complete listing of plant species and their distribution across the OHWM gradient. Some species occur in more than one category depending on site conditions. For example Indian plum and red alder may straddle the OHWM where soil drainage is high. They may occur above OHWM where soil drainage is low to moderate.

	Soil and geomorphic indicators <sup>24</sup>	Vegetative indicators <sup>25</sup>	Other indicators
At or straddling OHWM	<ul style="list-style-type: none"> <li>○ Top of bank</li> <li>○ Toe of lowest terrace (if terrace has developed horizons which may include a duff layer and A and B horizons versus freshly deposited alluvium)</li> <li>○ Benches</li> </ul>	<ul style="list-style-type: none"> <li>○ Willows</li> <li>○ Western red cedar</li> <li>○ Vine maple (streams)</li> <li>○ Black cottonwood</li> <li>○ Red alder <i>reed</i></li> <li>○ Salmonberry <i>canary grass</i></li> <li>○ Nootka rose</li> <li>○ Maidenhair and lady fern</li> <li>○ Blackberries</li> <li>○ Dunegrasses</li> </ul>	<ul style="list-style-type: none"> <li>☒ Sediment lines on vegetation or other fixed objects</li> <li>○ Change from channel deposits to older alluvium.</li> <li>☒ Darker stain lines on fixed objects</li> <li>○ Exposed roots/root scour.</li> <li>○ Drainage patterns, as evidenced by flattened vegetation</li> <li>○ Weathered and buried driftwood</li> </ul>
Above OHWM	<ul style="list-style-type: none"> <li>○ Hillslope toe</li> <li>○ Terraces or alluvium with an organic horizon or other developed soil horizons</li> <li>☒ Relic floodplain surface</li> <li>☒ Well developed soil A and B horizons/duff layer</li> </ul>	<ul style="list-style-type: none"> <li>○ Indian plum</li> <li>○ Red alder</li> <li>○ Western red cedar</li> <li>○ Douglas fir</li> <li>○ Western hemlock</li> <li>○ Ponderosa pine</li> <li>☒ Oregon white oak</li> <li>○ Coast pine</li> <li>○ Quaking aspen</li> <li>○ Vine maple (lakes)</li> <li>☒ Blackberries</li> </ul>	<ul style="list-style-type: none"> <li>☒ Lighter or no staining on fixed objects</li> <li>○ Overbank deposits</li> </ul>

### Notes

At this site sediment deposits on cobble and riprap were used, along with wrack on the sandy parts of the shoreline and the lowest extent of some native sedges.



Plant Distribution Across OHWM Gradient					
Below <del>Above</del> OHWM		At/Straddling OHWM		Above OHWM	
large cobble or riprap		reed canarygrass	few	black cottonwood	NI
				Oregon white oak	FACU
				Himalaya blackberry	FACU

20 Nov 2019  
Photo Direction: SW  
WSEL: 7.24ft NAVD88

LR-B-(1-6)



OHWM: Sediment line on rocks

# Appendix A: Field data form

## General Information

Site/Project: Wapato Valley /  
 Name/Owner: Plas Newydd Farm  
 Location: Lewis River  
 Description: 45.800006, -122.762723  
points: LR-C-(1-3)

The following field form is for use in the field to help in making ordinary high water mark delineations on streams. The form should be used as a guide. A team consisting of a hydrologist/ geomorphologist and a biologist may be needed to accurately determine the ordinary high water mark.

## General Observations: Day of Site Visit


Date of site visit:	11-19-2019		
Time of site visit:	15:35		
Weather conditions:	Overcast		
Watershed development:	Highly developed <input checked="" type="radio"/>	Mod. Developed <input type="radio"/>	Undeveloped <input type="radio"/>
Reach development:	Highly developed <input checked="" type="radio"/>	Mod. Developed <input type="radio"/>	Undeveloped <input type="radio"/>
Recent site disturbance?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe:
Upstream flow control devices?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: Merwin Dam
Bank armoring at the site?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: opposite (north) shoreline
Bank armoring up or downstream?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: north shoreline
Observable tidal backwater?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	
In-water structures? (i.e. bridge pilings, railroad embankments)	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: Railroad bridge, wood habitat structures, pilings
Animals grazing in riparian zone?	No <input checked="" type="radio"/>	Yes <input type="radio"/>	Describe:
Observable beaver activity?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: fresh beaver chewed logs

## Complete Vegetation Transects

- Use guidelines in Chapter 4 to complete vegetation transects.
- Determine upper and lower bounds of the OHWM from vegetation transects.
- After completing vegetation transects, look for more field indicators near the upper and lower bounds of the OHWM. Use the checklist as guidance.

## Sketch

If a simple site, sketch a cross-sectional diagram of the site below. Include location of the waterway and upper and lower bounds of the OHWM defined by the vegetation communities or other OHWM indicators. Page 3 of the data form can be used for more complex sketches



## Additional Indicators

Check the indicators that are observable at the site that provide rationale for establishing the OHWM at this location. The rationale should be described in detail in the report and should be supported with photographs taken during the site visit.

	Soil and geomorphic indicators <sup>24</sup>	Vegetative indicators <sup>25</sup>	Other indicators
Below OHWM	<ul style="list-style-type: none"> <li><input type="checkbox"/> Sediment bars</li> <li><input checked="" type="checkbox"/> Scour line</li> <li><input type="checkbox"/> Clean cobbles/boulders.</li> <li><input checked="" type="checkbox"/> Bank erosion/scour</li> <li><input checked="" type="checkbox"/> Lack of soil horizons</li> </ul>	Vegetation tolerant of inundation or high flow disturbances such as: <ul style="list-style-type: none"> <li><input type="checkbox"/> Willows</li> <li><input type="checkbox"/> Black cottonwood</li> <li><input type="checkbox"/> Japanese knotweed</li> <li><input type="checkbox"/> Skunk cabbage</li> <li><input checked="" type="checkbox"/> Aquatic plants</li> </ul>	<ul style="list-style-type: none"> <li><input checked="" type="checkbox"/> Exposed roots/root scour</li> <li><input type="checkbox"/> Drainage patterns, as shown by flattened vegetation.</li> <li><input checked="" type="checkbox"/> Aquatic animals</li> <li><input type="checkbox"/> Algal mats</li> <li><input type="checkbox"/> Iron staining</li> </ul>

<sup>24</sup> Refer to Chapter 4 for a more complete description of indicators.


<sup>25</sup> Species are provided as examples. Refer to Appendix B for a more complete listing of plant species and their distribution across the OHWM gradient. Some species occur in more than one category depending on site conditions. For example Indian plum and red alder may straddle the OHWM where soil drainage is high. They may occur above OHWM where soil drainage is low to moderate.



	Soil and geomorphic indicators <sup>24</sup>	Vegetative indicators <sup>25</sup>	Other indicators
At or straddling OHWM	<ul style="list-style-type: none"> <li>○ Top of bank</li> <li>☒ Toe of lowest terrace (if terrace has developed horizons which may include a duff layer and A and B horizons versus freshly deposited alluvium)</li> <li>☒ Benches</li> </ul>	<ul style="list-style-type: none"> <li>○ Willows</li> <li>○ Western red cedar</li> <li>○ Vine maple (streams)</li> <li>○ Black cottonwood</li> <li>○ Red alder ☒ Red osier dogwood</li> <li>○ Salmonberry</li> <li>○ Nootka rose</li> <li>○ Maidenhair and lady fern</li> <li>○ Blackberries</li> <li>○ Dunegrasses</li> </ul>	<ul style="list-style-type: none"> <li>☒ Sediment lines on vegetation or other fixed objects</li> <li>○ Change from channel deposits to older alluvium.</li> <li>☒ Darker stain lines on fixed objects</li> <li>☒ Exposed roots/root scour.</li> <li>○ Drainage patterns, as evidenced by flattened vegetation</li> <li>☒ Weathered and buried driftwood</li> </ul>
Above OHWM	<ul style="list-style-type: none"> <li>☒ Hillslope toe</li> <li>☒ Terraces or alluvium with an organic horizon or other developed soil horizons</li> <li>☒ Relic floodplain surface</li> <li>☒ Well developed soil A and B horizons/duff layer</li> </ul>	<ul style="list-style-type: none"> <li>○ Indian plum ☒ Oregon ash</li> <li>○ Red alder</li> <li>○ Western red cedar</li> <li>○ Douglas fir ☒ Black cottonwood</li> <li>○ Western hemlock</li> <li>○ Ponderosa pine</li> <li>○ Oregon white oak</li> <li>○ Coast pine</li> <li>○ Quaking aspen</li> <li>○ Vine maple (lakes)</li> <li>☒ Blackberries</li> </ul>	<ul style="list-style-type: none"> <li>☒ Lighter or no staining on fixed objects</li> <li>○ Overbank deposits</li> </ul>

### Notes

This survey site was along the Lewis shoveline at the inlet of a side channel. OHWM indicators included bank erosion lines and wrack buildup along the base of the eroded shore.



Plant Distribution Across OHWM Gradient					
Below <del>Above</del> OHWM		At/Straddling OHWM		Above OHWM	
weed	canarygrass	PACW	weed	canarygrass	PACW
				himalaya blackberry	PACW
				black cottonwood	N1

19 Nov 2019

Photo Direction: SW

WSEL: 7.34ft NAVD88

LR-C-(1-3)



OHWM: Wrack buildup and benches



## Appendix A: Field data form

### General Information

Site/Project: Wapato Valley /  
 Name/Owner: Plas Newydd Farm  
 Location: Lewis River  
 Description: 45.85777, -122.769953  
points: LR-D-(1-7)

The following field form is for use in the field to help in making ordinary high water mark delineations on streams. The form should be used as a guide. A team consisting of a hydrologist/ geomorphologist and a biologist may be needed to accurately determine the ordinary high water mark.

### General Observations: Day of Site Visit


Date of site visit:	11-19-2019		
Time of site visit:	15:00		
Weather conditions:	overcast		
Watershed development:	Highly developed <input checked="" type="radio"/>	Mod. Developed <input type="radio"/>	Undeveloped <input type="radio"/>
Reach development:	Highly developed <input checked="" type="radio"/>	Mod. Developed <input type="radio"/>	Undeveloped <input type="radio"/>
Recent site disturbance?	No <input checked="" type="radio"/>	Yes <input type="radio"/>	Describe:
Upstream flow control devices?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: Merwin Dam / Bonneville Dam
Bank armoring at the site?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: on opposite (north) shoreline.
Bank armoring up or downstream?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: upstream both sides
Observable tidal backwater?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	
In-water structures? (i.e. bridge pilings, railroad embankments)	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: Railroad bridge, wood habitat structures, pilings
Animals grazing in riparian zone?	No <input checked="" type="radio"/>	Yes <input type="radio"/>	Describe:
Observable beaver activity?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: Beaver chewed sticks

### Complete Vegetation Transects

- Use guidelines in Chapter 4 to complete vegetation transects.
- Determine upper and lower bounds of the OHWM from vegetation transects.
- After completing vegetation transects, look for more field indicators near the upper and lower bounds of the OHWM. Use the checklist as guidance.

## Sketch

If a simple site, sketch a cross-sectional diagram of the site below. Include location of the waterway and upper and lower bounds of the OHWM defined by the vegetation communities or other OHWM indicators. Page 3 of the data form can be used for more complex sketches



## Additional Indicators

Check the indicators that are observable at the site that provide rationale for establishing the OHWM at this location. The rationale should be described in detail in the report and should be supported with photographs taken during the site visit.

	Soil and geomorphic indicators <sup>24</sup>	Vegetative indicators <sup>25</sup>	Other indicators
<b>Below OHWM</b>	<ul style="list-style-type: none"> <li><input type="checkbox"/> Sediment bars</li> <li><input type="checkbox"/> Scour line</li> <li><input type="checkbox"/> Clean cobbles/boulders.</li> <li><input checked="" type="checkbox"/> Bank erosion/scour</li> <li><input checked="" type="checkbox"/> Lack of soil horizons</li> </ul>	Vegetation tolerant of inundation or high flow disturbances such as: <ul style="list-style-type: none"> <li><input type="checkbox"/> Willows</li> <li><input type="checkbox"/> Black cottonwood</li> <li><input type="checkbox"/> Japanese knotweed</li> <li><input type="checkbox"/> Skunk cabbage</li> <li><input checked="" type="checkbox"/> Aquatic plants</li> </ul>	<ul style="list-style-type: none"> <li><input checked="" type="checkbox"/> Exposed roots/root scour</li> <li><input type="checkbox"/> Drainage patterns, as shown by flattened vegetation.</li> <li><input checked="" type="checkbox"/> Aquatic animals</li> <li><input type="checkbox"/> Algal mats</li> <li><input type="checkbox"/> Iron staining</li> </ul>

<sup>24</sup> Refer to Chapter 4 for a more complete description of indicators.


<sup>25</sup> Species are provided as examples. Refer to Appendix B for a more complete listing of plant species and their distribution across the OHWM gradient. Some species occur in more than one category depending on site conditions. For example Indian plum and red alder may straddle the OHWM where soil drainage is high. They may occur above OHWM where soil drainage is low to moderate.

	Soil and geomorphic indicators <sup>24</sup>	Vegetative indicators <sup>25</sup>	Other indicators
At or straddling OHWM	<ul style="list-style-type: none"> <li>○ Top of bank</li> <li>☒ Toe of lowest terrace (if terrace has developed horizons which may include a duff layer and A and B horizons versus freshly deposited alluvium)</li> <li>☒ Benches</li> </ul>	<ul style="list-style-type: none"> <li>○ Willows</li> <li>○ Western red cedar</li> <li>○ Vine maple (streams)</li> <li>○ Black cottonwood</li> <li>○ Red alder <i>reed</i></li> <li>○ Salmonberry <i>canary grass</i></li> <li>○ Noofka rose</li> <li>○ Maidenhair and lady fern</li> <li>○ Blackberries <i>western</i></li> <li>○ Dunegrasses <i>golden top</i></li> </ul>	<ul style="list-style-type: none"> <li>☒ Sediment lines on vegetation or other fixed objects</li> <li>☒ Change from channel deposits to older alluvium.</li> <li>☒ Darker stain lines on fixed objects</li> <li>☒ Exposed roots/root scour.</li> <li>○ Drainage patterns, as evidenced by flattened vegetation</li> <li>☒ Weathered and buried driftwood</li> </ul>
Above OHWM	<ul style="list-style-type: none"> <li>○ Hillslope toe</li> <li>☒ Terraces or alluvium with an organic horizon or other developed soil horizons</li> <li>☒ Relic floodplain surface</li> <li>☒ Well developed soil A and B horizons/duff layer</li> </ul>	<ul style="list-style-type: none"> <li>○ Indian plum <i>Oregon</i></li> <li>○ Red alder <i>Ash</i></li> <li>○ Western red cedar</li> <li>○ Douglas fir</li> <li>○ Western hemlock <i>Black</i></li> <li>○ Ponderosa pine <i>Hornwood</i></li> <li>○ Oregon white oak</li> <li>○ Coast pine</li> <li>○ Quaking aspen</li> <li>○ Vine maple (lakes)</li> <li>○ Blackberries</li> </ul>	<ul style="list-style-type: none"> <li>☒ Lighter or no staining on fixed objects</li> <li>○ Overbank deposits</li> </ul>

→ wrack deposited on sand

### Notes

The best OHWM indicator at this site was where there was a break due to erosion between the sandy shoreline and native sedges.



Plant Distribution Across OHWM Gradient					
Below <del>Above</del> OHWM		At/Straddling OHWM		Above OHWM	
reed canarygrass	FACW	reed canarygrass	FACW	bentgrass sp.	FAC
western golden-top	FACW	western golden-top	FACW	western golden-top	FACW
		slough sedge	ORSL	sand bar willow	FACW
				sneezeweed	FACW
				Oregon ash	FACW
				black cottonwood	NI
				sheep sorrel	FACW

19 Nov 2019

Photo Direction: NE  
WSEL: 7.57ft NAVD88

LR-D-(1-7)

OHWM: Erosion and wrack buildup





# Appendix A: Field data form

## General Information

Site/Project: Wapato Valley  
 Name/Owner: Plas Newydd Farm  
 Location: Lewis River  
 Description: 45.857695, -122.770411  
points: LR-E-(1-7)

The following field form is for use in the field to help in making ordinary high water mark delineations on streams. The form should be used as a guide. A team consisting of a hydrologist/ geomorphologist and a biologist may be needed to accurately determine the ordinary high water mark.

## General Observations: Day of Site Visit


Date of site visit:	NOV 19 2019		
Time of site visit:	15:13		
Weather conditions:	Overcast		
Watershed development:	Highly developed <input checked="" type="radio"/>	Mod. Developed <input type="radio"/>	Undeveloped <input type="radio"/>
Reach development:	Highly developed <input checked="" type="radio"/>	Mod. Developed <input type="radio"/>	Undeveloped <input type="radio"/>
Recent site disturbance?	No <input checked="" type="radio"/>	Yes <input type="radio"/>	Describe:
Upstream flow control devices?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: Merwin Dam / Bonneville Dam
Bank armoring at the site?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: on opposite (north) shoreline
Bank armoring up or downstream?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: upstream both sides
Observable tidal backwater?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	
In-water structures? (i.e. bridge pilings, railroad embankments)	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: railroad bridge, wood habitat structures, piling
Animals grazing in riparian zone?	No <input checked="" type="radio"/>	Yes <input type="radio"/>	Describe:
Observable beaver activity?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: Beavers chewed sticks

## Complete Vegetation Transects

- Use guidelines in Chapter 4 to complete vegetation transects.
- Determine upper and lower bounds of the OHWM from vegetation transects.
- After completing vegetation transects, look for more field indicators near the upper and lower bounds of the OHWM. Use the checklist as guidance.

## Sketch

If a simple site, sketch a cross-sectional diagram of the site below. Include location of the waterway and upper and lower bounds of the OHWM defined by the vegetation communities or other OHWM indicators. Page 3 of the data form can be used for more complex sketches



## Additional Indicators

Check the indicators that are observable at the site that provide rationale for establishing the OHWM at this location. The rationale should be described in detail in the report and should be supported with photographs taken during the site visit.

	Soil and geomorphic indicators <sup>24</sup>	Vegetative indicators <sup>25</sup>	Other indicators
Below OHWM	<ul style="list-style-type: none"> <li><input checked="" type="checkbox"/> Sediment bars</li> <li><input checked="" type="checkbox"/> Scour line</li> <li><input type="checkbox"/> Clean cobbles/boulders.</li> <li><input checked="" type="checkbox"/> Bank erosion/scour</li> <li><input checked="" type="checkbox"/> Lack of soil horizons</li> </ul>	Vegetation tolerant of inundation or high flow disturbances such as: <ul style="list-style-type: none"> <li><input type="checkbox"/> Willows</li> <li><input type="checkbox"/> Black cottonwood</li> <li><input type="checkbox"/> Japanese knotweed</li> <li><input type="checkbox"/> Skunk cabbage</li> <li><input checked="" type="checkbox"/> Aquatic plants</li> </ul>	<ul style="list-style-type: none"> <li><input checked="" type="checkbox"/> Exposed roots/root scour</li> <li><input type="checkbox"/> Drainage patterns, as shown by flattened vegetation.</li> <li><input checked="" type="checkbox"/> Aquatic animals</li> <li><input type="checkbox"/> Algal mats</li> <li><input type="checkbox"/> Iron staining</li> </ul>


<sup>24</sup> Refer to Chapter 4 for a more complete description of indicators.

<sup>25</sup> Species are provided as examples. Refer to Appendix B for a more complete listing of plant species and their distribution across the OHWM gradient. Some species occur in more than one category depending on site conditions. For example Indian plum and red alder may straddle the OHWM where soil drainage is high. They may occur above OHWM where soil drainage is low to moderate.

	Soil and geomorphic indicators <sup>24</sup>	Vegetative indicators <sup>25</sup>	Other indicators
At or straddling OHWM	<ul style="list-style-type: none"> <li>○ Top of bank</li> <li>☒ Toe of lowest terrace (if terrace has developed horizons which may include a duff layer and A and B horizons versus freshly deposited alluvium)</li> <li>☒ Benches</li> </ul>	<ul style="list-style-type: none"> <li>○ Willows</li> <li>○ Western red cedar</li> <li>○ Vine maple (streams)</li> <li>○ Black cottonwood</li> <li>○ Red alder</li> <li>○ Salmonberry <i>reed canopy</i></li> <li>○ Nootka rose <i>grass</i></li> <li>○ Maidenhair and lady fern</li> <li>○ Blackberries</li> <li>○ Dune grasses <i>goldenrods</i></li> </ul>	<ul style="list-style-type: none"> <li>☒ Sediment lines on vegetation or other fixed objects</li> <li>☒ Change from channel deposits to older alluvium.</li> <li>☒ Darker stain lines on fixed objects</li> <li>☒ Exposed roots/root scour.</li> <li>○ Drainage patterns, as evidenced by flattened vegetation</li> <li>☒ Weathered and buried driftwood</li> </ul>
Above OHWM	<ul style="list-style-type: none"> <li>○ Hillslope toe</li> <li>☒ Terraces or alluvium with an organic horizon or other developed soil horizons</li> <li>☒ Relic floodplain surface</li> <li>☒ Well developed soil A and B horizons/duff layer</li> </ul>	<ul style="list-style-type: none"> <li>○ Indian plum</li> <li>○ Red alder</li> <li>○ Western red cedar</li> <li>○ Douglas fir <i>willow</i></li> <li>○ Western hemlock</li> <li>○ Ponderosa pine</li> <li>○ Oregon white oak</li> <li>○ Coast pine <i>oregon ash</i></li> <li>○ Quaking aspen</li> <li>○ Vine maple (lakes)</li> <li>○ Blackberries</li> </ul>	<ul style="list-style-type: none"> <li>○ Lighter or no staining on fixed objects</li> <li>○ Overbank deposits</li> </ul>

Notes

OHWM indicators at this site included dark water stains on a large log and erosion along the river shoreline.




Plant Distribution Across OHWM Gradient					
Below <del>Above</del> OHWM		At/Straddling OHWM		Above OHWM	
reed canarygrass	FACW	sheep sorrel	FACU	sand bar willows	FACW
		bentgrass spp.	FAC	Oregon ash	FACW
		red-osier dogwood	FACW	Himalayan blackberry	FACU
		spirea	FACW	black cottonwood	
		western goldenrod	FACW	dovefoot geranium	
		reed canarygrass	FACW	slough sedge	OBL
		slough sedge	OBL		

19 Nov 2019

Photo Direction: SW

WSEL: 7.42ft NAVD88

LR-E-(1-7)



OHWM: Dark water stain on log and sediment erosion

## Appendix A: Field data form

### General Information

Site/Project: Wapato Valley /  
 Name/Owner: Plas Newydd Farm  
 Location: Lewis River  
 Description: 45.85357, -122.776643  
points: LR-F-(1-3)

The following field form is for use in the field to help in making ordinary high water mark delineations on streams. The form should be used as a guide. A team consisting of a hydrologist/ geomorphologist and a biologist may be needed to accurately determine the ordinary high water mark.

### General Observations: Day of Site Visit


Date of site visit:	20 NOV 2019		
Time of site visit:	10:42		
Weather conditions:	Full Sun		
Watershed development:	Highly developed <input checked="" type="radio"/>	Mod. Developed <input type="radio"/>	Undeveloped <input type="radio"/>
Reach development:	Highly developed <input checked="" type="radio"/>	Mod. Developed <input type="radio"/>	Undeveloped <input type="radio"/>
Recent site disturbance?	No <input checked="" type="radio"/>	Yes <input type="radio"/>	Describe:
Upstream flow control devices?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: Merwin Dam / Bonneville Dam
Bank armoring at the site?	No <input checked="" type="radio"/>	Yes <input type="radio"/>	Describe:
Bank armoring up or downstream?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: upstream
Observable tidal backwater?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	
In-water structures? (i.e. bridge pilings, railroad embankments)	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: Piling upstream
Animals grazing in riparian zone?	No <input checked="" type="radio"/>	Yes <input type="radio"/>	Describe:
Observable beaver activity?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: Beaver chews

### Complete Vegetation Transects

- Use guidelines in Chapter 4 to complete vegetation transects.
- Determine upper and lower bounds of the OHWM from vegetation transects.
- After completing vegetation transects, look for more field indicators near the upper and lower bounds of the OHWM. Use the checklist as guidance.

## Sketch

If a simple site, sketch a cross-sectional diagram of the site below. Include location of the waterway and upper and lower bounds of the OHWM defined by the vegetation communities or other OHWM indicators. Page 3 of the data form can be used for more complex sketches



## Additional Indicators

Check the indicators that are observable at the site that provide rationale for establishing the OHWM at this location. The rationale should be described in detail in the report and should be supported with photographs taken during the site visit.

	Soil and geomorphic indicators <sup>24</sup>	Vegetative indicators <sup>25</sup>	Other indicators
<b>Below OHWM</b>	<ul style="list-style-type: none"> <li>○ Sediment bars</li> <li>○ Scour line</li> <li>○ Clean cobbles/boulders.</li> <li><input checked="" type="checkbox"/> Bank erosion/scour</li> <li><input checked="" type="checkbox"/> Lack of soil horizons</li> </ul>	Vegetation tolerant of inundation or high flow disturbances such as: <ul style="list-style-type: none"> <li>○ Willows</li> <li>○ Black cottonwood</li> <li>○ Japanese knotweed</li> <li>○ Skunk cabbage</li> <li><input checked="" type="checkbox"/> Aquatic plants</li> </ul>	<ul style="list-style-type: none"> <li><input checked="" type="checkbox"/> Exposed roots/root scour</li> <li>○ Drainage patterns, as shown by flattened vegetation.</li> <li><input checked="" type="checkbox"/> Aquatic animals</li> <li>○ Algal mats</li> <li>○ Iron staining</li> </ul>

<sup>24</sup> Refer to Chapter 4 for a more complete description of indicators.


<sup>25</sup> Species are provided as examples. Refer to Appendix B for a more complete listing of plant species and their distribution across the OHWM gradient. Some species occur in more than one category depending on site conditions. For example Indian plum and red alder may straddle the OHWM where soil drainage is high. They may occur above OHWM where soil drainage is low to moderate.

	Soil and geomorphic indicators <sup>24</sup>	Vegetative indicators <sup>25</sup>	Other indicators
At or straddling OHWM	<ul style="list-style-type: none"> <li>○ Top of bank</li> <li>○ Toe of lowest terrace (if terrace has developed horizons which may include a duff layer and A and B horizons versus freshly deposited alluvium)</li> <li>✗ Benches</li> </ul>	<ul style="list-style-type: none"> <li>✗ Willows</li> <li>○ Western red cedar</li> <li>○ Vine maple (streams)</li> <li>○ Black cottonwood</li> <li>○ Red alder</li> <li>○ Salmonberry</li> <li>○ Nootka rose</li> <li>○ Maidenhair and lady fern</li> <li>○ Blackberries <i>reed canary grass</i></li> <li>○ Dunegrasses</li> </ul>	<ul style="list-style-type: none"> <li>○ Sediment lines on vegetation or other fixed objects</li> <li>✗ Change from channel deposits to older alluvium.</li> <li>○ Darker stain lines on fixed objects</li> <li>✗ Exposed roots/root scour.</li> <li>○ Drainage patterns, as evidenced by flattened vegetation</li> <li>✗ Weathered and buried driftwood</li> </ul>
Above OHWM	<ul style="list-style-type: none"> <li>○ Hillslope toe</li> <li>✗ Terraces or alluvium with an organic horizon or other developed soil horizons</li> <li>✗ Relic floodplain surface</li> <li>✗ Well developed soil A and B horizons/duff layer</li> </ul>	<ul style="list-style-type: none"> <li>○ Indian plum</li> <li>○ Red alder</li> <li>○ Western red cedar</li> <li>○ Douglas fir <i>willow</i></li> <li>○ Western hemlock</li> <li>○ Ponderosa pine <i>oregon ash</i></li> <li>○ Oregon white oak</li> <li>○ Coast pine <i>black cottonwood</i></li> <li>○ Quaking aspen</li> <li>○ Vine maple (lakes)</li> <li>○ Blackberries</li> </ul>	<ul style="list-style-type: none"> <li>✗ Lighter or no staining on fixed objects</li> <li>○ Overbank deposits</li> </ul>

### Notes

This survey site was at the confluence of the Lewis and Columbia Rivers. The OHWM here was not clear on any fixed objects so we used the lowest elevations of the native willows and slough sedge. Reed canarygrass was present but tolerates wet and dry conditions so it was not a useful indicator species.





Plant Distribution Across OHWM Gradient					
Below <del>Above</del> OHWM		At/Straddling OHWM		Above OHWM	
Canadian waterweed	OBL	slough sedge	OBL	willow	FACW
		willows	FACW	black cottonwood	NI
				rough cocklebur	FAC

20 Nov 2019  
Photo Direction: N  
WSEL: 8.42ft NAVD88

LR-F-(1-3)

OHWM: Start of sedge and willow species



## Appendix A: Field data form

### General Information

Site/Project: Wapato Valley /  
 Name/Owner: Plas Newydd Farm  
 Location: Lewis River  
 Description: 45.85255, -122.777056  
points: LR-G<sub>1</sub>-(1-2)

The following field form is for use in the field to help in making ordinary high water mark delineations on streams. The form should be used as a guide. A team consisting of a hydrologist/ geomorphologist and a biologist may be needed to accurately determine the ordinary high water mark.

### General Observations: Day of Site Visit


Date of site visit:	11-20-2019		
Time of site visit:	10:55		
Weather conditions:	Full sun		
Watershed development:	Highly developed <input type="radio"/>	Mod. Developed <input checked="" type="radio"/>	Undeveloped <input type="radio"/>
Reach development:	Highly developed <input checked="" type="radio"/>	Mod. Developed <input type="radio"/>	Undeveloped <input type="radio"/>
Recent site disturbance?	No <input checked="" type="radio"/>	Yes <input type="radio"/>	Describe:
Upstream flow control devices?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: Merwin and Bonneville Dam
Bank armoring at the site?	No <input checked="" type="radio"/>	Yes <input type="radio"/>	Describe:
Bank armoring up or downstream?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: upstream on both sides
Observable tidal backwater?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	
In-water structures? (i.e. bridge pilings, railroad embankments)	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: pilings upstream
Animals grazing in riparian zone?	No <input checked="" type="radio"/>	Yes <input type="radio"/>	Describe:
Observable beaver activity?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: fresh chewed sticks

### Complete Vegetation Transects

- Use guidelines in Chapter 4 to complete vegetation transects.
- Determine upper and lower bounds of the OHWM from vegetation transects.
- After completing vegetation transects, look for more field indicators near the upper and lower bounds of the OHWM. Use the checklist as guidance.

## Sketch

If a simple site, sketch a cross-sectional diagram of the site below. Include location of the waterway and upper and lower bounds of the OHWM defined by the vegetation communities or other OHWM indicators. Page 3 of the data form can be used for more complex sketches



## Additional Indicators

Check the indicators that are observable at the site that provide rationale for establishing the OHWM at this location. The rationale should be described in detail in the report and should be supported with photographs taken during the site visit.

	Soil and geomorphic indicators <sup>24</sup>	Vegetative indicators <sup>25</sup>	Other indicators
<b>Below OHWM</b>	<ul style="list-style-type: none"> <li><input checked="" type="checkbox"/> Sediment bars</li> <li><input type="checkbox"/> Scour line</li> <li><input type="checkbox"/> Clean cobbles/boulders.</li> <li><input checked="" type="checkbox"/> Bank erosion/scour</li> <li><input checked="" type="checkbox"/> Lack of soil horizons</li> </ul>	Vegetation tolerant of inundation or high flow disturbances such as: <ul style="list-style-type: none"> <li><input type="checkbox"/> Willows</li> <li><input type="checkbox"/> Black cottonwood</li> <li><input type="checkbox"/> Japanese knotweed</li> <li><input type="checkbox"/> Skunk cabbage</li> <li><input checked="" type="checkbox"/> Aquatic plants</li> </ul>	<ul style="list-style-type: none"> <li><input type="checkbox"/> Exposed roots/root scour</li> <li><input type="checkbox"/> Drainage patterns, as shown by flattened vegetation.</li> <li><input checked="" type="checkbox"/> Aquatic animals</li> <li><input type="checkbox"/> Algal mats</li> <li><input type="checkbox"/> Iron staining</li> </ul>


<sup>24</sup> Refer to Chapter 4 for a more complete description of indicators.

<sup>25</sup> Species are provided as examples. Refer to Appendix B for a more complete listing of plant species and their distribution across the OHWM gradient. Some species occur in more than one category depending on site conditions. For example Indian plum and red alder may straddle the OHWM where soil drainage is high. They may occur above OHWM where soil drainage is low to moderate.

	Soil and geomorphic indicators <sup>24</sup>	Vegetative indicators <sup>25</sup>	Other indicators
At or straddling OHWM	<ul style="list-style-type: none"> <li>○ Top of bank</li> <li>○ Toe of lowest terrace (if terrace has developed horizons which may include a duff layer and A and B horizons versus freshly deposited alluvium)</li> <li>○ Benches</li> </ul>	<ul style="list-style-type: none"> <li>⊗ Willows <i>sprouts</i></li> <li>○ Western red cedar</li> <li>○ Vine maple (streams)</li> <li>○ Black cottonwood</li> <li>○ Red alder <i>&amp; reed</i></li> <li>○ Salmonberry <i>canopy</i></li> <li>○ Nootka rose</li> <li>○ Maidenhair and lady fern</li> <li>○ Blackberries <i>blow in sedge</i></li> <li>○ Dune grasses</li> </ul>	<ul style="list-style-type: none"> <li>⊗ Sediment lines on vegetation or other fixed objects</li> <li>○ Change from channel deposits to older alluvium.</li> <li>⊗ Darker stain lines on fixed objects</li> <li>⊗ Exposed roots/root scour.</li> <li>○ Drainage patterns, as evidenced by flattened vegetation</li> <li>⊗ Weathered and buried driftwood</li> </ul>
Above OHWM	<ul style="list-style-type: none"> <li>○ Hillslope toe</li> <li>⊗ Terraces or alluvium with an organic horizon or other developed soil horizons</li> <li>⊗ Relic floodplain surface</li> <li>⊗ Well developed soil A and B horizons/duff layer</li> </ul>	<ul style="list-style-type: none"> <li>○ Indian plum <i>&amp; willows</i></li> <li>○ Red alder <i>(larger)</i></li> <li>○ Western red cedar <i>mature</i></li> <li>○ Douglas fir</li> <li>○ Western hemlock</li> <li>○ Ponderosa pine</li> <li>○ Oregon white oak</li> <li>○ Coast pine</li> <li>○ Quaking aspen</li> <li>○ Vine maple (lakes)</li> <li>○ Blackberries</li> </ul>	<ul style="list-style-type: none"> <li>○ Lighter or no staining on fixed objects</li> <li>⊗ Overbank deposits</li> </ul>

### Notes

This survey site was at the confluence of the Lewis and Columbia Rivers. Water levels here are highly variable as evidenced by lines of washed up debris/wrack at multiple elevations on the beach. Elevations were taken at larger wrack lines and where willows were sprouting a little higher on the beach.



Plant Distribution Across OHWM Gradient					
Below OHWM		At/Straddling OHWM		Above OHWM	
non-vegetated sand		sand bar willow	FACW	sitka willow	FACW
Eurasian watermilfoil	OBL	reed canarygrass	FACW	sand bar willow	FACW
		slough sedge	OBL	reed canarygrass	FACW
		rough cocklebur	FAC	Oregon ash	FACW
		western golden-top	FACW	black cottonwood	not listed
				western golden-top	FACW
				Robert geranium	FACW

20 Nov 2019  
Photo Direction: S  
WSEL: 8.42ft NAVD88

LR-G-(1-2)



OHWL: Change from sand to vegetation such as sedge and willow species

# Appendix A: Field data form

## General Information

Site/Project	Wapata Valley
Name/Owner:	Plas Newydd Farm
Location:	Lewis River
Description:	45.85581, -122.773755 Points LR-H-(1-5)

The following field form is for use in the field to help in making ordinary high water mark delineations on streams. The form should be used as a guide. A team consisting of a hydrologist/ geomorphologist and a biologist may be needed to accurately determine the ordinary high water mark.

## General Observations: Day of Site Visit

Date of site visit:	3 Dec 2019		
Time of site visit:	10:21		
Weather conditions:	Overcast		
Watershed development:	Highly developed <input checked="" type="radio"/>	Mod. Developed <input type="radio"/>	Undeveloped <input type="radio"/>
Reach development:	Highly developed <input checked="" type="radio"/>	Mod. Developed <input type="radio"/>	Undeveloped <input type="radio"/>
Recent site disturbance?	No <input checked="" type="radio"/>	Yes <input type="radio"/>	Describe:
Upstream flow control devices?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: BONIVUL
Bank armoring at the site?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: across the river
Bank armoring up or downstream?	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe:
Observable tidal backwater?	No <input type="radio"/>	Yes <input type="radio"/>	
In-water structures? (i.e. bridge pilings, railroad embankments)	No <input type="radio"/>	Yes <input checked="" type="radio"/>	Describe: boat dock on other side of river
Animals grazing in riparian zone?	No <input checked="" type="radio"/>	Yes <input type="radio"/>	Describe:
Observable beaver activity?	No <input checked="" type="radio"/>	Yes <input type="radio"/>	Describe:


## Complete Vegetation Transects

- o Use guidelines in Chapter 4 to complete vegetation transects.
- o Determine upper and lower bounds of the OHWM from vegetation transects.
- o After completing vegetation transects, look for more field indicators near the upper and lower bounds of the OHWM. Use the checklist as guidance.



## Sketch

If a simple site, sketch a cross-sectional diagram of the site below. Include location of the waterway and upper and lower bounds of the OHWM defined by the vegetation communities or other OHWM indicators. Page 3 of the data form can be used for more complex sketches



## Additional Indicators

Check the indicators that are observable at the site that provide rationale for establishing the OHWM at this location. The rationale should be described in detail in the report and should be supported with photographs taken during the site visit.

	Soil and geomorphic indicators <sup>24</sup>	Vegetative indicators <sup>25</sup>	Other indicators
<b>Below OHWM</b>	<input checked="" type="checkbox"/> Sediment bars <input type="checkbox"/> Scour line <input type="checkbox"/> Clean cobbles/boulders. <input checked="" type="checkbox"/> Bank erosion/scour <input checked="" type="checkbox"/> Lack of soil horizons	Vegetation tolerant of inundation or high flow disturbances such as: <input type="checkbox"/> Willows <input type="checkbox"/> Black cottonwood <input type="checkbox"/> Japanese knotweed <input type="checkbox"/> Skunk cabbage <input checked="" type="checkbox"/> Aquatic plants	<input type="checkbox"/> Exposed roots/root scour <input checked="" type="checkbox"/> Drainage patterns, as shown by flattened vegetation. <input checked="" type="checkbox"/> Aquatic animals <input type="checkbox"/> Algal mats <input type="checkbox"/> Iron staining

reed canary grass

<sup>24</sup> Refer to Chapter 4 for a more complete description of indicators.


<sup>25</sup> Species are provided as examples. Refer to Appendix B for a more complete listing of plant species and their distribution across the OHWM gradient. Some species occur in more than one category depending on site conditions. For example Indian plum and red alder may straddle the OHWM where soil drainage is high. They may occur above OHWM where soil drainage is low to moderate.

	Soil and geomorphic indicators <sup>24</sup>	Vegetative indicators <sup>25</sup>	Other indicators
At or straddling OHWM	<ul style="list-style-type: none"> <li>○ Top of bank</li> <li>☒ Toe of lowest terrace (if terrace has developed horizons which may include a duff layer and A and B horizons versus freshly deposited alluvium)</li> <li>☒ Benches</li> </ul>	<ul style="list-style-type: none"> <li>○ Willows</li> <li>○ Western red cedar</li> <li>○ Vine maple (streams)</li> <li>○ Black cottonwood</li> <li>○ Red alder</li> <li>○ Salmonberry</li> <li>○ Nootka rose</li> <li>○ Maidenhair and lady fern</li> <li>○ Blackberries</li> <li>○ Dunegrasses</li> </ul>	<ul style="list-style-type: none"> <li>○ Sediment lines on vegetation or other fixed objects</li> <li>○ Change from channel deposits to older alluvium.</li> <li>○ Darker stain lines on fixed objects</li> <li>○ Exposed roots/root scour.</li> <li>○ Drainage patterns, as evidenced by flattened vegetation</li> <li>☒ Weathered and buried driftwood</li> </ul>
Above OHWM	<ul style="list-style-type: none"> <li>○ Hillslope toe</li> <li>☒ Terraces or alluvium with an organic horizon or other developed soil horizons</li> <li>☒ Relic floodplain surface</li> <li>☒ Well developed soil A and B horizons/duff layer</li> </ul>	<ul style="list-style-type: none"> <li>○ Indian plum</li> <li>○ Red alder</li> <li>○ Western red cedar</li> <li>○ Douglas fir</li> <li>○ Western hemlock</li> <li>○ Ponderosa pine</li> <li>○ Oregon white oak</li> <li>○ Coast pine</li> <li>○ Quaking aspen</li> <li>○ Vine maple (lakes)</li> <li>○ Blackberries</li> </ul>	<ul style="list-style-type: none"> <li>○ Lighter or no staining on fixed objects</li> <li>○ Overbank deposits</li> </ul>

Oregon ash and  
Black cotton wood

### Notes

There is a wrack line that is an OHWM indicator.



Plant Distribution Across OHWM Gradient					
Below					
Above OHWM		At/Straddling OHWM		Above OHWM	
Canadian waterweed	OBL	reed canarygrass		FACW reed canarygrass	FACW
reed canarygrass	FACW	slough sedge		OBL Oregon ash	FACW
needle spikerush	OBL			sand bar willow	FACW
				Pacific willow	FACW
				bentgrass sp.	FAC

3 Dec 2019

Photo Direction: NE

WSEL: 8.31ft NAVD88

LR-H-(1-5)

OHWM: Wrack line



---

**ATTACHMENT B**

**SPECIES AND COMMON NAMES OF PLANTS**

---

<b>Common Camas</b>	<b>Species Name</b>
Bird's Foot Trefoil	<i>Lotus corniculatus</i>
Black Cottonwood	<i>Populus balsamifera trichocarpa</i>
Black Hawthorn	<i>Crataegus douglasii</i>
Bur-reed	<i>Sparganium sp</i>
Common Camas	<i>Camassia quamish</i>
Douglas Fir	<i>Pseudotsuga douglasii</i>
Douglas Spirea	<i>Spirea douglasii</i>
False Indigo Bush	<i>Amorpha fruticosa</i>
Herb Robert	<i>Geranium robertianum</i>
Himalayan Blackberry	<i>Rubus armeniascus</i>
Licorice Fern	<i>Polypodium glycerrhiza</i>
Needle Spikerush	<i>Eleocharis acicularis</i>
Oregon Ash	<i>Fraxinus latifolia</i>
Oregon White Oak	<i>Quercus garryana</i>
Red-Osier Dogwood	<i>Cornus alba</i>
Reed Canarygrass	<i>Phalaris arundinacea</i>
Rough Cocklebur	<i>Xanthium strumarium</i>
Scot's Broom	<i>Cystisus scoparius</i>
Slough Sedge	<i>Carex obnupta</i>
Smartweed	<i>Polygonum sp</i>
Snowberry	<i>Symphoricarpos albus</i>
Softstem Bulrush	<i>Schoenoplectus tabernaemontanii</i>
Wapato	<i>Sagittaria latifolia</i>
Western Goldenrod	<i>Euthamia occidentalis</i>
Willows	<i>Salix sp</i>
Woolgrass	<i>Scirpus cyperinus</i>
Wormleaf Stonecrop	<i>Sedum stenopelatum</i>

**From:** [Kelley Jorgensen](#)  
**To:** [Brent Davis](#); [Jenna Kay](#)  
**Subject:** [Contains External Hyperlinks] FW: Zip shapes for delivery  
**Date:** Wednesday, December 4, 2019 4:58:14 PM  
**Attachments:** [image003.png](#)  
[100yr Flood zone PN Farm Property.zip](#)  
[Wetland Rating Units Wapato.zip](#)  
[OHWM PN FARM & Wapato Bank.zip](#)

**CAUTION:** This email originated from outside of Clark County. Do not click links or open attachments unless you recognize the sender and know the content is safe.

Hi Brent and Jenna,

Please find attached the wetland rating unit, OHWM and 100-year flood GIS shapefiles for the Plas Newydd property.

These are the locations we are proposing updates for the Shoreline Master Plan process.

OHWM delineation technical memo to follow under separate cover.

Please don't hesitate to contact me or Chris Watson if you have any questions.

Thank you for your time and consideration,

Kelley



---

**KELLEY JORGENSEN** » *President of Conservation*


*she | her | hers*

T 360.857.4087 c 971.285.6874 E [kjorgensen@pnfarm.com](mailto:kjorgensen@pnfarm.com)

PO Box 428 | Ridgefield, WA 98642 | [www.wapato-valley.com](http://www.wapato-valley.com)

Please find attached GIS shapefiles:

- 100yr flood zone on the PN Farm property based on the USACE 1% exceedance value at Columbia RM 87 of 26.54' NAVD 88.



- Wetland rating units on the Wapato Mitigation and Conservation Bank provided by CEG.
- OHWM from determination.

---

**CHRIS WATSON** » *GIS Manager/Project*

*Manager/Geologist*

*he | him | his*

T 360.857.4040 c 503.332.7231 E [cwatson@pnfarm.com](mailto:cwatson@pnfarm.com)



---

PO Box 428 | Ridgefield, WA 98642 | [www.wapato-valley.com](http://www.wapato-valley.com)



**From:** [David Morgan](#)  
**To:** [Davis, Brent](#)  
**Cc:** [Kay, Jenna](#); [Pool, Bob](#); [Kelley Jorgensen](#)  
**Subject:** [Contains External Hyperlinks] RE: Shoreline Review  
**Date:** Thursday, November 7, 2019 1:45:24 PM

---

**CAUTION:** This email originated from outside of Clark County. Do not click links or open attachments unless you recognize the sender and know the content is safe.

Brent,

Thank you.

We are working on a shapefile for your team to review and should have that in the middle of next week if that is soon enough for your timeline. For the IRT review you mention below lets chat next week about what you need there. I will ask Chris to work directly with Bob on the GIS file types and details.

Thank you very much.

David



---

**DAVID MORGAN** » *Managing Partner*

*he | him | his*

T 360.857.4039 C 503.750.7570 E [dmorgan@pnfarm.com](mailto:dmorgan@pnfarm.com)

PO Box 428 | Ridgefield, WA 98642 | [www.wapato-valley.com](http://www.wapato-valley.com)

---

**From:** Davis, Brent <[Brent.Davis@clark.wa.gov](mailto:Brent.Davis@clark.wa.gov)>  
**Sent:** Tuesday, November 5, 2019 10:09 AM  
**To:** David Morgan <[dmorgan@pnfarm.com](mailto:dmorgan@pnfarm.com)>  
**Cc:** Kay, Jenna <[Jenna.Kay@clark.wa.gov](mailto:Jenna.Kay@clark.wa.gov)>; Pool, Bob <[ROBERT.POOL@clark.wa.gov](mailto:ROBERT.POOL@clark.wa.gov)>  
**Subject:** RE: Shoreline Review

David,

The Shoreline Map is adopted with the Shoreline Master Program and can only be changed through a Shoreline Amendment process, so now is the time. I suggest you contact Bob Pool to get the data formatting details for our GIS and then submit your OHWM (pursuant to RCW 90.58), Wetland, and Floodplain mapping to Jenna for inclusion in the upcoming map amendment. It would also be helpful to have concurrence documentation from the IRT for the record.



**Brent Davis**  
Wetland and Habitat Review Manager

COMMUNITY DEVELOPMENT

564.397.4152



---

**From:** Kay, Jenna  
**Sent:** Tuesday, November 5, 2019 9:54 AM  
**To:** 'David Morgan'  
**Cc:** Davis, Brent  
**Subject:** RE: Shoreline Review

Hi David,

Thank you for reaching out. I am going to connect you with Brent Davis, copied here, the county's Shoreline Administrator. His team will be able to follow-up with you further.

Regards,  
Jenna



**Jenna Kay**  
Planner II  
COMMUNITY PLANNING

564.397.4968



---

**From:** David Morgan [mailto:dmorgan@pnfarm.com]  
**Sent:** Monday, November 4, 2019 4:57 PM  
**To:** Kay, Jenna  
**Subject:** [Contains External Hyperlinks] Shoreline Review

**CAUTION:** This email originated from outside of Clark County. Do not click links or open attachments unless you recognize the sender and know the content is safe.

Jenna,

I would like to setup a time to talk with the department about some inconsistencies I see in the proposed mapping for my property in North West Clark County along the Lewis River. We are working with Clark County on a mitigation and conservation bank and it looks like the Ordinary High Water line on my property does not reflect the work we have done to date. If you can let me know the best way to come in and have a conversation about this I would appreciate it.

Thank you  
David Morgan

**DAVID MORGAN** » *Managing Partner*

---

T 360.857.4039 C 503.750.7570 E dmorgan@pnfarm.com  
PO Box 428 | Ridgefield, WA 98642 | [www.wapato-valley.com](http://www.wapato-valley.com)

This e-mail and related attachments and any response may be subject to public disclosure under state law.